

## **Three new *Hieracium* taxa (*Asteraceae*, *Cichorieae*) from Mt Cholomon, Chalkidiki, Central Makedonia, Greece**

Author(s): Günter Gottschlich, Konstantina Melikoki, Eleni Eleftheriadou & Konstantinos Theodoropoulos

Source: Willdenowia, 43(1):59-64.

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

<https://doi.org/10.3372/wi.43.43106>

URL: <http://www.bioone.org/doi/full/10.3372/wi.43.43106>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

GÜNTER GOTTSCHLICH<sup>1</sup>\*, KONSTANTINA MELIKOKI<sup>2</sup>, ELENI ELEFThERiADOU<sup>2</sup> & KONSTANTINOS THEODOROPOULOS<sup>2</sup>

## Three new *Hieracium* taxa (*Asteraceae*, *Cichorieae*) from Mt Cholomon, Chalkidiki, Central Makedonia, Greece

### Abstract

Gottschlich G., Melikoki K., Eleftheriadou E. & Theodoropoulos K.: Three new *Hieracium* taxa (*Asteraceae*, *Cichorieae*) from Mt Cholomon, Chalkidiki, Central Makedonia, Greece. – Willdenowia 43: 59–64. June 2013. – Online ISSN 1868-6397; © 2013 BGBM Berlin-Dahlem.

Stable URL: <http://dx.doi.org/10.3372/wi.43.43106>

Three new subspecies, *Hieracium sparsum* subsp. *cholomonense*, *H. transiens* subsp. *levimaculatum* and *H. bohatschianum* subsp. *onosmoidiforme*, all from Mt Cholomon, Chalkidiki, Central Makedonia, Greece, are described and illustrated.

Additional key words: local endemic, taxonomy, *Compositae*

### Introduction

The flora of Chalkidiki area was known from earlier floristic and vegetation studies (Ganiatsas 1963; Dafis 1966; Voliotis 1967; Pavlides 1976; Athanasiadis & Theodoropoulos 1990; Theodoropoulos 1991; Athanasiadis & al. 1992; Babalonas & al. 1998; Damianidis 2011; Karousou & al. 2008). However, because of many other interesting plants which can be found in Greece, the genus *Hieracium* L. (*Asteraceae*, *Cichorieae*) has often not attracted necessary interest by field botanists in the past. Within a project of floristic investigation of Mt Cholomon, the Greek co-authors collected with special emphasis on *Hieracium*, because it is known that intensive botanical investigation of restricted areas often reveals local taxa, especially in apomictic genera such as *Hieracium*. The specimens collected were revised by G. Gottschlich. Among them three taxa proved to be undescribed.

### Results and Discussion

**1. *Hieracium sparsum* subsp. *cholomonense* Gottschl. & Melikoki, **subsp. nov.** – Fig. 1.**

Holotype: Greece, Central Makedonia, Prefecture of Chalkidiki, Mt Cholomon, 40°26'14.7"N, 23°30'18.4"E, *Quercus frainetto* forest, 890 m, 6 Jul 2011, K. Melikoki 12.4.2 (TAUF; isotypes: B, C, herb. Gottschlich 57781). Paratypes: Mt Cholomon, 40°26'0.9"N, 23°30'20.2"E, *Quercus frainetto* forest, 860 m, 1 Jul 2010, K. Melikoki 4.2.1 (TAUF); Mt Cholomon, 40°26'8.67"N, 23°30'14.96"E, slope of *Quercus frainetto* forest, 852 m, 6 Jul 2011, K. Melikoki 12.1.4 (TAUF); Mt Cholomon, 40°26'9.1"N, 23°29'55.7"E, *Quercus frainetto* forest, 816 m, 6 Jul 2011, K. Melikoki 13.3.3 (TAUF); Mt Cholomon, 40°26'14.6"N, 23°30'16.1"E, *Quercus frainetto* forest, 879 m, 6 Jul 2011, K. Melikoki 12.3.2 (TAUF); Mt Cholomon, 40°26'4.1"N, 23°30'13.7"E, *Quercus frainetto*

1 Hermann-Kurz-Str. 35, 72074 Tübingen, Germany; \*email: ggtuebingen@yahoo.com (author for correspondence).

2 Institute of Forest Botany-Geobotany, Faculty of Forestry and Natural Environment, Aristotle University of Thessaloniki, GR-54124, Thessaloniki, Greece; email: melikoki@gmail.com; eelefthe@for.auth.gr; ktheodor@for.auth.gr

forest, 833 m, 6 Jul 2011, K. Melikoki, 13.5.5 (TAUF); Mt Cholomon, 40°26'37"N, 23°29'32.6"E, *Quercus frainetto* forest, 686 m, 9 Jul 2011, K. Melikoki 15.2.1 (TAUF, herb. Gottschlich 57780).

*Description* — *Planta* perennis. *Rhizoma* tenue vel crassum, horizontale vel obliquum. *Caulis* erectus, solidus, (25–)30–60(–70) cm altus, infra medium rubro-vel brunneo-viridis (ad basim saepe rubro-violaceus), supra viridis, subtiliter striatus, phyllopodus, pilis simplicibus infra sparsis, supra nullis vel solitariis, 1–1.5 mm longis, mollibus, albis, denticulatis, pilis glanduliferis nullis, pilis stellatis inferne sparsis, superne nullis. *Folia rosularia* (3 vel)4 vel 5(vel 6), petiolata; *petiolus* 1–2 cm longus, rubro-violaceus, pilis simplicibus subdensis vel densis, 1.5–2 mm longis, mollibus, albis, denticulatis, pilis glanduliferis perparvulis sparsis, pilis stellatis nullis; *lamina* late lanceolato-obovata vel elliptica, subpapyracea, 3–7 × 1.5–3 cm, obscure viridis vel olivacea, interdum pauce et indistincte brunneo-maculata, margine remote denticulata vel breviter et acute dentata, ad basim longe attenuata, apice acuta, supra calvescens vel pilis simplicibus sparsis, in marginem et subtus pilis modice densis, in costa dorsali subdensis vel densis, 1.5–2 mm longis, albis, mollibus, denticulatis, pilis glanduliferis perparvulis tantum in margine et in costa dorsali sparsis, pilis stellatisque nullis. *Folia caulina* 1 vel 2(vel 3), inferior lanceolato-obovatum, lamina petioliformiter attenuata, denticulata, reliqua lanceolato-lineata, sessilia, indumento nullo. *Synflorescentia* laxo paniculata; *rami* (3 vel)4–8(–10), stricti, tenues, 2–15 cm longi, quisque calathidiis 1–3(–5); *calathidia* (3–)5–20(–30); acladium 1.5–2 cm longum. *Pedunculus* tenuissimus (tantum <0.5 mm diametro), indumento nullo, sub involucrio 2–5 bracteis dilute viridibus, basibus alatis, 1.5–2 mm longis. *Involucrum* 9–10 mm longum, anguste campanulatum. *Involucra phylla* pauciseriata,



Fig. 1. *Hieracium sparsum* subsp. *cholomonense* – A: habit; B: indumentum of involucral bracts. – Isotype: K. Melikoki 12.4.2 (herb. Gottschlich 57781).

exteriora 2–5 mm longa, obscure olivacea, immarginata, ovata vel anguste ovata, subobtusata, interiora 9–10 mm longa, olivacea, pallide marginata, lanceolata, ad basim circa 1.5 mm lata, acuta, pilis simplicibus sparsis, 1 mm longis, mollibus, ad basim atris, partibus reliquis albis, pilis glanduliferis sparsis vel modice densis, 0.2–0.6 mm longis, capite luteis, pilis stellatis nullis. *Ligulae* linguatae, luteae, glabrae. *Styli* lutei, papillis nigris. *Areolae* margine dentatae. *Achaenia* 3–3.1 mm longa, atrobrunnea. *Florescentia*: Iulio mense.

*Taxonomy* — As Szelağ (2003) pointed out, “the *Hieracium sparsum* group (*H. sparsum* Friv. s.l. sensu Zahn

Table 1. Comparison of diagnostic characters of *Hieracium sparsum* subsp. *cholomonense*, *H. sparsum* subsp. *macedonicum* and *H. sparsum* subsp. *borbasii*.

Diagnostic character	<i>Hieracium sparsum</i> subsp. <i>cholomonense</i>	<i>Hieracium sparsum</i> subsp. <i>macedonicum</i>	<i>Hieracium sparsum</i> subsp. <i>borbasii</i>
Number of cauline leaves	1 or 2(or 3)	3–7	4–8
Peduncles			
stellate hairs	absent	few	moderate
glandular hairs	absent	absent	few to numerous
simple hairs	absent	absent	few to numerous
Involucre [mm]	9–10	9–12	10–12
Involucral bracts			
apex	acute	obtuse	subacute
stellate hairs	absent	absent	absent
glandular hairs	few to numerous	few to numerous	few to numerous
simple hairs	few	few to numerous	few to numerous
Distribution	N Greece	Albania, Republic of Macedonia, N Greece	S Carpathians: Retezat Mts <sup>1</sup>

<sup>1</sup>Zahn (1922–1938) mentioned also a record from Bulgaria (as *Hieracium sparsum* var. *latifolium*), but it must be proved whether or not this taxon belongs to *H. sparsum* subsp. *borbasii*. Szélag (2006) mentioned the same uncertain case concerning a record of *H. sparsum* subsp. *borbasii* from the Pojana Rusca Mts in Romania.

1921–1923) constitutes a separate and morphologically fairly uniform complex of mountain, especially sub-alpine, taxa occurring in SE Europe and W Asia". In Europe there exist two diversity centres: the Balkan Peninsula and the Southern Carpathians. Taxa described during the 19th century mostly were established at the species level. Later, Zahn (1921–1923, 1922–1938) treated them at the rank of subspecies within one collective species. Alone or together with his local contributors he described numerous further taxa at the rank of subspecies or even variety, subvariety or form. Recently Szélag (2003), who generally favours species rank within the *H. sparsum* complex, listed in his "Synopsis of *Hieracium* sect. *Cernua* (*Asteraceae*)" 43 species and 30 subspecies. In his "Taxonomic revision of *Hieracium* sect. *Cernua* (*Asteraceae*) in the Carpathians, Sudetes and Alps" (Szélag 2006) he dealt with all taxa at species rank. In the Med-Checklist (Greuter 2008) the Central European taxonomic concept with collective species and subspecies was still maintained. In the ongoing Checklist for Greece this concept will also be used. Although morphological features already hypothesize that there may exist taxa of different value within the *H. sparsum* complex, we accept here for practical reasons, i.e. working on a steady base, the concept of the Med-Checklist. Future molecular analyses may bring more light into this complex. For a first study, dealing with two taxa of *H. sect. Cernua* R. Uechtr., see Ronikier & Szélag (2008).

To classify taxa of *Hieracium sparsum* some important features must be considered: shape and indumentum of basal leaves; number, shape, insertion and indumentum of stem leaves; and ramification, size and indumentum of capitula. Considering these features, the new *H.*

*sparsum* subsp. *cholomonense* is closely related to *H. sparsum* subsp. *borbasii* (R. Uechtr.) Zahn and *H. sparsum* subsp. *macedonicum* (Boiss. & Orph.) Zahn. The diagnostic characters of these three taxa are compared in Table 1.

## 2. *Hieracium transiens* subsp. *levimaculatum* Gottschl. & Melikoki, **subsp. nov.** – Fig. 2.

Holotype: Greece, Central Makedonia, Prefecture of Chalkidiki, Mt Cholomon, 40°25'56.6"N, 23°33'34.5"E, *Quercus frainetto* forest, 751 m, 8 Jun 2011, K. Melikoki 7.3.3 (TAUF; isotypes: B, C, herb. Gottschlich 57839).

Paratypes: Mt Cholomon, 40°26'8.8"N, 23°32'18.3"E, *Quercus frainetto* forest, 778 m, 8 Jun 2011, K. Melikoki 7.11.1 (ATHU, B, C, TAUF, W, herb. Gottschlich 57779).

*Description* — *Planta* perennis. *Rhizoma* tenue vel crassum, horizontale vel obliquum. *Caulis* erectus, solidus, (35–)40–50(–60) cm altus, infra medium rubro- vel brunneo-viridis (ad basim saepe rubro-violaceus), supra viridis, subtiliter striatus, phyllopodus, pilis simplicibus infra sparsis, supra sparsis vel nullis, 1–1.5 mm longis, mollibus, albis, denticulatis, pilis glanduliferis perparvulis (circa 0.05 mm tantum) sparsis, pilis stellatis nullis vel sparsis. *Folia rosularia* (3 vel)4 vel 5(–7), petiolata; *petiolus* (1.5–)2–3(–3.5) cm longus, rubro-violaceus, pilis simplicibus subdensis vel densis, 1.5–2 mm longis, mollibus, albis, denticulatis, pilis glanduliferis perparvulis sparsis, pilis stellatis nullis; *lamina* anguste elliptica vel late lanceolata, 3–6 × 1.5–2 cm, supra plusminusve glauco-viridia vel olivacea, disperse vel modice brunneo-maculata, subtus rubro-violacea, margine denticulata vel

breviter et acute dentata, ad basim cuneata, apice acuta, pilis simplicibus sparsis, in marginem et subtus in costa dorsali subdensis, 1–1.5 mm longis, albis, mollibus, denticulatis, pilis glanduliferis stellatisque nullis. *Folia caulina* 1 vel 2, inferior ut in foliis rosularibus, petiolatum, reliquum multo angustius, saepe lineariformiter reductum. *Synflorescentia* laxe paniculata; *rami* 3–5 (vel 6), stricti vel arcuato-ascendentes, tenues, (3–)5–10(–25) cm longi, quisque calathidiis 1–3(–6); *calathidia* (4 vel)5–8(–10); *accladium* 1–1.3 cm longum. *Pedunculus* tenuis, pilis simplicibus sparsis vel modice densis, atris (apice albis), 0.5–1 mm longis, pilis glanduliferis sparsis vel modice densis, 0.3–0.5 mm longis, capite obscure luteis, pilis stellatis sparsis vel subdensis, sub involucri 1 vel 2 bracteis atro-viridibus, 1.5–2 mm longis. *Involucrum* 9–10 mm longum, anguste campanulatum. *Involucris phylla* pauciseriata, exteriora 4–5 mm longa, obscure olivacea, anguste marginata, anguste ovata, subobtusata, interiora 9–10 mm longa, dilute viridia, lanceolata, ad basim 1–1.1 mm lata, acuta, pilis simplicibus subdensis, circa 1 mm longis, atris (apice albis), pilis glanduliferis sparsis, 0.2–0.3 mm longis, capite luteis, pilis stellatis in marginem subdensis, apice barbulata. *Ligulae* lingulatae, luteae, glabrae. *Styli* lutei. *Areolae* margine fibrilloso-dentatae. *Achaenia* circa 3 mm longa, obscure brunnea. *Florescentia*: Iunio et Iulio mense.

*Taxonomy* — *Hieracium transiens* is morphologically intermediate between *H. sparsum* and *H. murorum* L. As a result of the great morphological variation of the presumptive parental species there exist numerous often locally endemic taxa. In shape and mottling of leaves and in indumentum there exist no hitherto described taxa that correspond with the new subspecies. The most similar is *H. transiens* subsp. *zygosense* (Zahn) Greuter, described from Thessalia (Zahn 1921–23: 1043), but that taxon has more lanceolate, unspotted leaves and long acute involucri bracts with fewer stellate hairs.



Fig. 2. *Hieracium transiens* subsp. *levimaculatum* – A: habit; B: indumentum of involucri bracts. – Isotype: K. Melikoki 7.3.3 (herb. Gottschlich 57839).

**3. *Hieracium bohatschianum* subsp. *onosmoidiforme* Gottschl. & Melikoki, subsp. nov.** – Fig. 3.

*Holotype*: Greece, Central Makedonia, Prefecture of Chalkidiki, Mt Cholomon, 40°30'13.6"N, 23°33'13.5"E, slope of road, rocky position, 547 m, 6 Jun 2011, K. Melikoki 6.12.3 (TAUF; isotypes: ATHU, B, C, FI, LD, PAL, UPA, W, herb. Gottschlich 57774).

*Paratype*: [Greece, Central Makedonia, Prefecture of Chalkidiki], Cholomon-Gebirge, zwischen Arnaea und der Passhöhe, Silikatschutthang, 800 m, 14 Jun 1979, A. Polatschek (W 1979-16038).

*Description* — *Planta* perennis. *Rhizoma* tenue vel crassum, horizontale vel obliquum. *Caulis* erectus, solidus, (20–)25–40(–50) cm altus, infra medium rubro- vel brunneo-viridis (ad basim saepe rubro-violaceus), supra viridis, subtiliter striatus, phyllopodus, pilis simplicibus infra subdensis, supra nullis, 3–5 mm longis, mollibus, albis, denticulatis, pilis glanduliferis nullis, pilis stellatis



Fig. 3. *Hieracium bohatschianum* subsp. *onosmoidiforme* – A: habit; B: indumentum of involucreal bracts. – Isotype: K. Melikoki 6.12.3 (herb. Gottschlich 57774).

sparsis. *Folia rosularia* (2 vel)3 vel 4(vel 5), petiolata; *petiolus* 1–2 cm longus, rubro-violaceus, pilis simplicibus subdensis vel densis, 3–5 mm longis, mollibus, albis, denticulatis, pilis glanduliferis perparvulis sparsis, pilis stellatis nullis; *lamina* late lanceolato-obovata vel elliptica, 5–8 × 2.5–3.5 cm, supra canescenti-viridis vel glauco-viridis, subtus rubro-violacea, margine integra vel denticulata, apice acuta, supra calvescens, versus marginem pilis simplicibus sparsis, subrigidis, in marginem modice densis, in costa dorsali densis, 2.5–4 mm longis, albis, submollibus, denticulatis, pilis glanduliferis perparvulis tantum in margine et in costa dorsali sparsis, pilis stellatis nullis. *Folia caulina* (2 vel)3(vel 4), inferior late

lanceolatum, reliqua anguste ovata, denticulata, sessilia vel subamplexicaulia, color et indumento ut in foliis basalibus. *Synflorescentia* laxe paniculata; *rami* 3–6, stricti, tenues, 2–5(–8) cm longi, quisque calathidiis 1 vel 2; *calathidia* (3–)5–7(–10); *acladium* 1–2 cm longum. *Pedunculus* tenuis, pilis simplicibus sparsis, 1.3–1.7 mm longis, mollibus, ad basim atris, partibus reliquis albis, pilis glanduliferis sparsis, 0.2–0.3 mm longis, capite luteis, pilis stellatis sparsis, sub involucre 1 vel 2 bracteis viridibus, 1.5–2 mm longis. *Involucrum* 9–10 mm longum, semi-ellipsoideum. *Involucri phylla* pauciseriata, obscure olivacea, late pallide marginata, lanceolata, ad basim 1–1.2 mm lata, acuta, pilis simplicibus sparsis, circa 1 mm longis, mollibus, ad basim atris, partibus reliquis albis, pilis glanduliferis modice densis, 0.2–0.3 mm longis, capite luteis, pilis stellatis sparsis. *Ligulae* lingulatae, luteae, glabrae. *Styli* lutei. *Areolae* margine breviter denticatae. *Achaenia* circa 3 mm longa, obscure brunnea. *Florescentia*: Iunio mense.

*Taxonomy* — *Hieracium bohatschianum* Zahn was originally described from Mt Trescovaț in Romania (Zahn 1910). Later another two subspecies, each only known from one locality in the Re-

public of Macedonia, were described: *H. bohatschianum* subsp. *platytrigonodon* O. Behr & al. from Mt Korab (Behr & al. 1937) and *H. bohatschianum* subsp. *heterodontophyes* O. Behr & al. (Behr & al. 1939). In 1989 the first author received for determination a specimen from A. Polatschek, Vienna, collected in 1979 on Mt Cholomon. Because of its intermediate characters we already then assumed that it must belong to *H. bohatschianum*. Not having seen the type specimen or the other described subspecies we added a “sensu lato” to the determination. In the meantime we could study type material of *H. bohatschianum* subsp. *platytrigonodon* issued within the exsiccata series “Herbarium Hieraciorum”, ed. Otto

Behr, Forst, Lusatiae No. 975 (B, BASBG, HBG, M). Recently Szelağ (2011) re-collected *H. bohatschianum* at the type locality. Because of the lack of a holotype in BP he selected a neotype and added an instructive figure to his publication. Szelağ confirmed the intermediate morphological position (“sparsum – schmidtii”) of *H. bohatschianum*. Comparing all this we came to the conclusion that the collection from Mt Cholomon is new to science. It more closely resembles the type specimen of *H. bohatschianum* than the two taxa described from the Republic of Macedonia, both of which have more elongated leaves and only one stem leaf; furthermore their leaves are more toothed. The newly described *H. bohatschianum* subsp. *onosmoidiforme* differs from *H. bohatschianum* in having basal leaves with distinct and longer petioles and stem leaves less amplexicaul.

## Acknowledgements

We thank Zbigniew Szelağ (Kraków) and an anonymous reviewer for valuable comments on the manuscript.

## References

- Athanasiadis N. & Theodoropoulos K. 1990: Contribution to the knowledge of the Mount Cholomon flora. – Sci. Ann. Dept. Forest. Nat. Environm. Aristot. Univ. Thessaloniki **32(2)**: 167–211. [In Greek with English summary].
- Athanasiadis N., Theodoropoulos K. & Konstantinidis P. 1992: The phytosociological units in the forests of *Pinus nigra* ssp. *pallasiana* in Sithonia Peninsula of Chalkidiki, N. Greece. – Sci. Ann. Dept. Forest. Nat. Environm. Aristot. Univ. Thessaloniki Vol. **35(1)**: 273–306. [In Greek with English summary].
- Babalonas D., Constantinou M. & Charalambides S. 1998: I chlorida tou Agiou Orous [The flora of Mount Athos]. – Thessaloniki: Organization Cultural Capital “Thessaloniki 1997”. [In Greek].
- Behr O., Behr E. & Zahn K. H. 1937: Beitrag zur Kenntnis der Hieracien der Balkanhalbinsel. – Glasn. Skopsk. Nauč. Društva **18**: 51–67.
- Behr O., Behr E. & Zahn K. H. 1939: Beiträge zur Kenntnis der Hieracien von Südserbien. – Glasn. Skopsk. Nauč. Društva **20**: 23–34.
- Dafis S. 1966: Standorts- und ertragskundliche Untersuchungen in Eichen- und Kastanienniederwälder der N.Ö. Chalkidiki. – Habilitation, Aristotle University of Thessaloniki. [in Greek with German summary].
- Damianidis C. 2011: Phytosociological research of the *Erica* heathlands and evergreen broadleaved shrublands of the north side of the mount Cholomon. – MSc Thesis, Faculty of Forestry and Natural Environment, Aristotle University of Thessaloniki. [In Greek with English summary].
- Ganiatsas C. A. 1963: I vlastisis ke I chloris tis chersonisou tou Agiou Orous [The vegetation and flora of the peninsula of Holy Mountain]. – Athoniki Politia: 509–678. Thessaloniki.
- Greuter W. 2008: *Dicotyledones (Compositae)*. – In: Greuter W. & Raab-Straube E. von (ed.), Med-Checklist. A critical inventory of vascular plants of the circum-mediterranean countries **2**. Genève: OPTIMA.
- Karousou R., Hanlidou E., Kokkini P., Koufou D. & Kokkini S. 2008: On the flora of Mount Stratonikon (GR1270005), a NATURA 2000 site of N Greece. – Fl. Medit. **18**: 529–550.
- Pavlidis G. 1976: On the flora and vegetation of the Sithonia peninsula (Chalkidiki, Greece). – PhD Thesis, Aristotle University of Thessaloniki. [In Greek with English summary].
- Ronikier M. & Szelağ Z. 2008: Differentiation among disjunct populations of agamospermous species of *Hieracium* section *Cernua* (Asteraceae) in Central European subalpine habitats. – Bot. J. Linn. Soc. **158**: 93–105.
- Szelağ Z. 2003: A synopsis of *Hieracium* sect. *Cernua* (Asteraceae). – Polish Bot. J. **48**: 89–97.
- Szelağ Z. 2006: Taxonomic revision of *Hieracium* sect. *Cernua* (Asteraceae) in the Carpathians, Sudets and Alps. – Polish Bot. J. **51**: 97–153.
- Szelağ Z. 2011: Hieracia Balcanica VII. Identity and typification of *Hieracium bohatschianum* (Asteraceae). – Polish Bot. J. **56**: 155–157.
- Theodoropoulos K. 1991: Bestimmung und Klassifizierung der pflanzensoziologischen Vegetationseinheiten im Universitätswald Taxiarchis Chalkidiki. – Sci. Ann. Fac. Agric. Forest. Univ. Thessaloniki **18**: 1–200 + tables. [In Greek with German abstract].
- Voliotis D. 1967: Untersuchungen über die Vegetation und die Flora des Cholomongebirges unter besonderer Berücksichtigung der aromatischen, Heil- und Bienezuchtpflanzen. – Sci. Ann. Fac. Phys. Math. Aristot. Univ. Thessaloniki **10**: 1–131. [In Greek with German summary].
- Zahn K. H. 1910: Die ungarischen Hieracien des National-Museums zu Budapest, zugleich V. Beitrag zur Kenntnis der Hieracien Ungarns und Balkanländer. – Ann. Hist.-Nat. Mus. Natl. Hung. **8**: 34–106.
- Zahn K. H. 1921–1923: *Hieracium*. – In: Engler A. (ed.), Das Pflanzenreich **75**: 1–288, **76**: 289–576, **77**: 577–864 (1921); **79**: 865–1146 (1922); **82**: 1147–1705 (1923). – Leipzig: Engelmann.
- Zahn K. H. 1922–1938: *Hieracium*. – In: Ascherson P. F. A. & Graebner K. O. P. P. (ed.), Synopsis der mitteleuropäischen Flora **12(1)**: 1–80 (1922), 81–160 (1924), 161–400 (1929), 401–492 (1930); **12(2)**: 1–160 (1930), 161–480 (1931), 481–640 (1934), 641–790 (1935); **12(3)**: 1–320 (1936), 321–480 (1937), 481–708 (1938). – Leipzig & Berlin: Borntraeger.