

Cryptogamie, Bryologie, 2008, 29 (2): 161-000
 © 2008 Adac. Tous droits réservés

Towards the bryophyte flora of Greece, studies in Chalkidiki area (North Greece)

Marko SABOVLJEVIĆ^a, Evdoxia TSAKIRI^b and Aneta SABOVLJEVIĆ^a

^a*Institute of Botany and Botanical Garden, Faculty of Biology,
 University of Belgrade, Takovska 43, 11000 Belgrade, Serbia*

^b*Laboratory of Systematic Botany and Phytogeography, Department of Botany,
 School of Biology, Aristotle University of Thessaloniki, Thessaloniki 541 24,
 Greece*

(Received 25 May 2007, accepted 27 July 2007)

Abstract – The bryophyte flora of the Chalkidiki Prefecture (*Nomos*) in Greece has been studied, yielding a record of 138 bryophytes, (22 liverwort and 116 mosses). Twenty three species are newly recorded for Chalkidiki, 18 for north-east floristic region of Greece, 3 for the Greek mainland and one for Greece.

Bryophytes / Flora / Distribution / Chalkidiki / Greece

Résumé – La flore bryophytique de la Préfecture Chalkidiki (Nomos) en Grèce a été étudiée. Au total, 138 bryophytes (22 hépatiques et 116 mousses) ont été censées. Vingt-trois espèces représentent de nouveaux registres pour Chalkidiki, 18 pour la région floristique du NE de la Grèce, 3 pour la Grèce continentale et une pour la Grèce.


Bryophytes / Flore / Distribution / Chalkidiki / Grèce

INTRODUCTION

Although Greece is bryologically one of the best known regions of South Eastern Europe, it is among the worst known regions of Southern Europe (Sabovljević, 2004). The knowledge of its bryophyte flora remains insufficient, and it is believed that there is still a lot to be found (ECCB, 1995; Düll, 1996; Söderström *et al.*, 1998; Sabovljević *et al.*, 2001). Most of the bryophyte records were due to foreign researchers who mostly visited Greece on holidays. Thus some islands, such as, *e.g.*, the island of Crete, are incomparably bryologically better known than the mainland of Greece.

Few bryophyte checklists have been published for Greece (Preston, 1981, 1984a-b; Düll, 1995a; Ros *et al.*, 2007). Additionally, new records and floristic contributions have been recently reported for the country (Mavrommatis, 1972;

* Correspondence and reprints: marko@worldpost.eu



Athanasiadis, 1977; Giannitsaros & Koumpli-Sovantzi, 1990; Blockeel, 1991, 2001; Turland & Wilson, 1995; Carratello & Aleffi, 1998; Gallego & Cano, 1998; Papp *et al.*, 1998a-b; Papp, 1998, 2002, 2003; Blockeel *et al.*, 2002; Maier, 2002; Lüth, 2002, 2003; Tsakiri & Babalonas, 2002; Lara *et al.*, 2003; Tsakiri *et al.*, 2005, 2006; Henderson & Blockeel, 2006; Blockeel & Ramel, 2006; Parent & De Zuttere, 2006).

The first publication concerning Chalkidiki was the “*Florae Graecae Prodromus*” (Sibthorp & Smith, 1806-1813) on a collection from Athos Mt. Since then, more data were published by Podpéra (1922), Ganiatsas (1938), Taylor (1952), Politis (1953), Reimers (1957), Vajda (1959), Geissler (1977), Zoller *et al.* (1977), Gamisans & Hébrard (1980), Düll (1995b), Sauer (1995), Papp & Babalonas (1996) and Erzberger (2006). Nowadays *ca* 252 taxa are mentioned to occur in Chalkidiki areas, among the *ca* 740 taxa reported so far from the country (*ca* 520 of them in North Greece), which suggests that much floristic and ecological investigation is still to be done in this area to define its bryophyte flora.

MATERIAL AND METHODS

Study area

The Chalkidiki peninsula is situated in the southern part of Central Macedonia District (North Greece). The peninsula has specific shape of a three-fingered hand spreading out in a north-south direction into the Aegean Sea, and is attached to the mainland of Macedonia at its northern part. Each of the three fingers of Chalkidiki peninsula consists in a small sub-peninsula named from west to east Kassandra, Sithonia and Athos (Fig. 1). The maximum elevations of the 3 sub-peninsulas increase from west to east: 353m in Kassandra, 817m in Sithonia, and 2033m in Athos.

Geologically, Chalkidiki is mainly basic with calcareous rocks, but other geological formations can be found around the peninsula, from sand to crystalline rocks, granodiorite or schists (Mountrakis, 1985). The climate is Eu-Mediterranean, with dry summers and most precipitation during winter period. Mean annual temperature varies from *ca* 15.9°C by the sea areas (e.g. Agios Mamas) to *ca* 12.7°C on the mountain (e.g. Arnea). The average annual precipitation varies accordingly between 500-800 mm (Balafoutis 1977). With the exception of urban and cultivated areas, the peninsula is mostly covered by *Pinus halepensis* forest with Mediterranean maquis, frequently dominated by *Erica* shrubs, with many areas in different degradation stages.

The easternmost sub-peninsula (Athos) is probably the most interesting area. The elevation of Athos Mt (2033m) together with the lack of urbanisation, form a unique environmental status that promises interesting findings. However, it is bryologically still unexplored, since the area south of the Ouranoupolis village, is closed for visitors (*Holly Athos Mt. Monastery Community*). The Natura 2000 sites within the study area are (areas at the sea are not included here): *Oros Cholomontas*, *Oros Itamos-Sithonia*, *Limnohalassa Agiou Mama*, *Oros Stratoniukon* (Chalkidiki Prefecture), and *Limnothalassa Epanomis*, *Limnothalassa Angelochoriou* (Thessaloniki Prefecture) on the way from Thessaloniki to Chalkidiki (Dafis *et al.*, 1996).

Field work was conducted at the Chalkidiki peninsula (excluding closed Athos area, south of Ouranoupolis) and southeastern area of Thessaloniki prefecture in late autumn during the years 2002-2006.



Fig. 1. North Greece and the location of the study area: Chalkidiki Prefecture and part of Thessaloniki Prefecture (the shady area at the smaller map of Greece). The smaller map shows also the Greek floristic regions (for abbreviations and their definitions see the text). Circles (●) indicate the collection sites. Altitudinal lines are at 600, 1,000 and 1,400m a.s.l.

Methodology

In total 46 localities with wider surroundings have been prospected, and collections were made focusing on all habitat types. The locality names are given in groups according to the sub-peninsula where they are situated. Most of the sites are included in Chalkidiki Prefecture, while only six sites (Nº: 2, 5, 6, 19, 43, 44) are situated in Thessaloniki Prefecture (Table 1, Fig. 1). The voucher specimens are kept at the Herbarium of the Belgrade University (BEOU).

The nomenclature for liverworts follows Sabovljević & Natcheva (2006), while that of mosses follows Hill *et al.* (2006), and for each taxon, the localities where it was recorded have been expressed by the corresponding numbers in Table 1.

RESULTS AND DISCUSSION

A total of 138 bryophyte taxa were recorded from the collection sites; 22 of which are liverworts, while 116 are mosses.

The taxa are listed alphabetically, first hepaticas and then mosses. For each taxon, data on the habitat (substratum), and locality numbers are given (for details, see Table 1 and Fig. 1).



Table 1. The investigated localities with geographical coordinates and altitude, sorted regionally and alphabetically (see also Fig.1).

	<i>Locality Nº</i>	<i>N</i>	<i>E</i>	<i>Altitude (m)</i>
Region of Thessaloniki				
Agia Paraskevi	2	40°28'53"	23°03'	230
East of Thessaloniki	5	40°38"	22°58'05"	ca 200
Epanomi	6	40°25'41"	22°55'45"	50
Nea Michaniona	19	40°28"	22°51'45"	30
South-East of Thessaloniki	43	40°30'30"	22°58'36"	10
Vasilika	44	40°28'53"	23°08'15"	75
Chalkidiki mainland & 3rd peninsula				
Agios Mamas	3	40°15'42"	23°20'	15
Gerakini	8	40°16'47"	23°26'53"	20
Ierissos	10	40°23'48"	23°52'34"	10
Nea Kallikratia	18	40°19"	23°03'56"	20
Nea Moudania	20	40°15'32"	23°17'	20
Nea Roda	22	40°22'51"	23°55'25"	10
Nea Triglia	24	40°18'24"	23°12'36"	90
Olynthos (antiq.)	27	40°17'05"	23°21'17"	40
Ormilia	28	40°17'42"	23°32'45"	40
Ouranoupoli	30	40°19'33"	23°59'	10
Petalona	33	40°22'06"	23°09'42"	270
Pirgadikia	38	40°20'20"	23°43'34"	20
Poligiro	34	40°23'37"	23°26'25"	530
Psakoudia	37	40°15'47"	23°29'28"	10
Vatopedi	45	40°15'34"	23°33'32"	20
Kassandra Peninsula (1st peninsula)				
Afitos	1	40°06'	23°26'10"	50
Fourka - village	7	40°00'21"	23°25'	40
Hanioti	9	40°00'05"	23°34'30"	10
Kalandra	12	39°58'18"	23°20'07"	60
Kalithea	13	40°04'13,5"	23°26'47"	50
Kassandria	14	40°03"	23°24'40"	50
Kriopigi	15	40°02'09"	23°28'21"	130
Nea Fokea	17	40°08"	23°23'49"	20
Nea Potidea	21	40°11'43,5"	23°19'38,5"	20
Nea Skioni	23	39°56'52"	23°32'	10
Paliouri	31	39°56'35"	23°39'49"	125
Pefkochori	32	39°59'17"	23°36'40"	20
Polichrono	35	40°01"	23°31'40"	20
Sani	39	40°14"	23°20'	5
Siviri	42	40°02'09"	23°21'41"	5
Sithonia Peninsula (2nd peninsula)				
Agios Nikolaos	4	40°15"	23°41'54"	100
Kalamitsi	11	39°59'19"	23°59'	15
Metamorfosi	16	40°14"	23°34'21"	20
Neos Marmaras	25	40°05'40"	23°47'08"	20
Nikiti	26	40°13'48"	23°40'21"	40
Ormos Panagias	29	40°13'48"	23°44'21"	10
Porto Carras	36	40°04'29"	23°47'30"	10
Sarti	40	40°05'50"	23°58'36"	10
Sikia	41	40°02'23"	23°57'	40
Vourvourou	46	40°11'11"	23°47'34"	20



The following symbols apply:

+ : taxa new to NE floristic region of Greece


(a) : taxa new to Chalkidiki (already reported for NE floristic region of Greece)

Liverworts


1. *Calypogeia fissa* (L.) Raddi – deep soil – 25, 30
2. + *Cephaloziella baumgartneri* Schiffn. – wet soil in slopes and cliffs – 1, 9, 13, 17, 23, 28, 38, 42
3. *Conocephalum conicum* (L.) Dumort. – wet rocks and soil – 2, 5, 13, 21, 22, 24, 27, 30, 33, 37, 41, 46
4. + *Conocephalum salebrosum* Szweykowski, Buczkowska & Odrzykoski – wet rocks-34
5. *Corsinia coriandrina* (Spreng.) Lindb. – soil – 2, 25, 37, 29, 41
6. *Frullania dilatata* (L.) Dumort. – tree bark – 3, 4, 15, 26, 36
7. *Frullania tamarisci* (L.) Dumort. – soil in macchia – 40
8. *Gongylanthus ericetorum* (Raddi) Nees – soil - 26
9. *Lejeunea cavifolia* (Ehrh.) Lindb. – wood – 27, 29, 46
10. *Lophocolea bidentata* (L.) Dumort. – soil - 25
11. *Lophocolea heterophylla* (Schrad.) Dumort. – wood – 25, 40, 45
12. *Lunularia cruciata* (L.) Lindb. – soil – 7, 10, 11
13. *Marchantia polymorpha* L. subsp. *ruderale* Bischl. & Boisselier – wet soil by the leaking water – 2, 13
14. *Metzgeria furcata* (L.) Dumort – schistous rocks – 34, 37
15. *Pellia endiviifolia* (Dicks.) Dumort. – rock and soil by the water – 2, 3, 18, 23
16. *Reboulia hemisphaerica* (L.) Raddi – soil – 6, 8, 19, 31, 33, 43
17. *Riccia ciliifera* Link ex Lindenb. – soil – 11, 16, 36
18. *Riccia nigrella* DC. – soil – 11, 36
19. *Riccia sorocarpa* Bisch. – soil – 40
20. *Scapania compacta* (A. Roth) Dumort. – soil – 25, 41
21. *Scapania undulata* (L.) Dumort. – soil - 25
22. *Southbya tophacea* (Spruce) Spruce – soil – 7, 26

Mosses


1. *Aloina aloides* (Koch ex Schulz) Kindb. – soil and slopes – 3, 7, 15, 17, 18, 20, 35, 44
2. *Aloina ambigua* (Bruch & Schimp.) Limpr. – soil – 10, 15, 24, 31, 32, 33, 36, 42, 44, 46
3. *Aloina rigida* (Hedw.) Limpr. – soil – 2, 4, 5, 6, 9, 16, 21, 30, 43, 46
4. *Amblystegium serpens* (Hedw.) Schimp. – rocks – 11, 16, 19, 22, 28
5. *Anomodon viticulosus* (Hedw.) Hook. & Taylor – tree base – 25, 45
6. *Atrichum undulatum* (Hedw.) P. Beauv. – soil – 10, 14, 25, 37,
7. *Barbula convoluta* Hedw. – soil – 4, 12, 33, 44
8. *Barbula commutata* Jur. – soil – 17, 21, 29, 39



9. *Barbula unguiculata* Hedw. – soil – all sites
10. *Bartramia stricta* Bruch & Schimp. – soil – 46
11. *Brachytheciastrum velutinum* (Hedw.) Ignatov & Huttunen – soil and tree bark – 11, 18, 30, 42, 46
12. *Bryoerythrophyllum recurvirostrum* (Hedw.) P.C. Chen – soil – 5, 8, 14, 19, 31
13. *Bryum argenteum* Hedw. – soil – all
14. + *Bryum argenteum* var. *lanatum* (P.Beauv.) Hampe – soil – 15, 24, 32, 39
15. + *Bryum canariense* Brid. – soil - 25
16. *Bryum capillare* Hedw. – soil – 3, 4, 10, 11, 13, 14, 15, 19, 30, 35, 36, 37, 43, 44
17. *Bryum dichotomum* Hedw. – soil – 11, 24, 26, 31
18. *Bryum moravicum* Podp. – wood – 8, 27, 39, 40, 41
19. *Bryum pseudotriquetrum* (Hedw.) P. Gaertn. et al. – soil by water – 25
20. *Campylophyllum calcareum* (Crundw. & Nyholm) Hedenäs – soil among rocks – 26, 37
21. *Campylopus oerstadianus* (Müll. Hal.) Mitt. – soil on schistous rocks – 25
22. *Ceratodon purpureus* (Hedw.) Brid. – soil – 25
23. + *Crossidium crassinerve* (De Not.) Jur. – soil and slopes – 15
24. *Crossidium squamiferum* (Viv.) Jur. – soil among rocks and boulders – 23
25. *Dicranella heteromalla* (Hedw.) Schimp. – soil - 27
26. *Dicranella howei* Renauld & Cardot – soil – 6, 12, 15, 36
27. *Dicranella varia* (Hedw.) Schimp. – soil – 7, 13
28. *Dicranum scoparium* Hedw. – soil - 10
29. *Didymodon acutus* (Brid.) K. Saito – soil – 33, 34
30. *Didymodon insulanus* (De Not.) M. O. Hill – soil – 8, 9, 14, 22, 38, 43
31. *Didymodon luridus* Hornsch. – soil – 1, 2, 3, 6, 14, 15, 16, 17, 19, 20, 21, 22, 28, 30, 35, 37, 42, 45, 46
32. *Didymodon rigidulus* Hedw. – soil – 4, 7, 9, 14, 16, 22, 27, 29 39, 41
33. + *Didymodon sicculus* M. J. Cano, Ros, Garcia-Zamora & J. Guerra –soil – 25, 31
34. + *Didymodon umbrosus* (Müll.Hal.) R. H. Zander –soil – 9, 15, 32
35. *Didymodon vinealis* (Brid.) K. Saito – soil – 1, 3, 4, 8, 12, 19, 22, 23, 31, 32, 42
36. *Encalypta streptocarpa* Hedw. – rocks and walls 33, 34
37. *Encalypta vulgaris* Hedw. – soil – 13, 39
38. *Eucladium verticillatum* (With.) Bruch & Schimp. – wet rocks – 13, 23
39. *Fissidens bryoides* Hedw. – soil - 31
40. *Fissidens dubius* P. Beauv. – soil – 16, 25
41. + *Fissidens serrulatus* Brid. – soil - 26
42. *Fissidens taxifolius* Hedw. – soil – 25
43. *Fontinalis antipyretica* Hedw. – in water – 25
44. *Funaria hygrometrica* Hedw. – on soil – 8, 14, 17, 19, 22, 39
45. + *Funariella curviseta* (Schwägr.) Sérgio – rocky wall – 16





46. *Grimmia decipiens* (Schultz) Lindb. – siliceous rock – 25, 26
47. + *Grimmia dissimulata* E. Maier – rock – 10
48. *Grimmia lisae* De Not. – siliceous rock – 25, 26, 29, 37
49. *Grimmia pulvinata* (Hedw.) Sm. – rocks – 4, 11, 12, 15, 28, 36, 42
50. *Grimmia trichophylla* Grev. – rocks – 9, 11, 32
51. *Gymnostomum calcareum* Nees & Hornsch. – wet rocks – 13
52. + *Gymnostomum lanceolatum* M. J. Cano, Ros & J. Guerra – shaded rock – 23
53. *Gyroweissia tenuis* (Hedw.) Schimp. – rock – 35, 42
54. *Hedwigia ciliata* (Hedw.) P. Beauv. – siliceous rock – 26
55. *Homalothecium aureum* (Spruce) H. Rob. – rocks – 27
56. *Homalothecium lutescens* (Hedw.) H. Rob. – rocks – 12, 18, 29
57. *Homalothecium sericeum* (Hedw.) Schimp. – rocks – 16, 21, 28, 37, 38
58. *Hypnum cupressiforme* Hedw. – tree bark and soil – 3, 6, 8, 11, 12, 17, 22, 32, 35, 40
59. *Isothecium alopecuroides* (Lam. ex Dubois.) Isov. – tree base – 46
60. *Isothecium myosuroides* Brid. – siliceous rock – 46
61. *Kindbergia praelonga* (Hedw.) Ochyra – stones and soil – 25, 26
62. *Leptodon smithii* (Hedw.) F. Weber & D. Mohr – tree trunk – 25, 34
63. *Leucodon sciurooides* (Hedw.) Schwägr. – tree trunk – 25
64. (a) *Microbryum starkeanum* (Hedw.) R. H. Zander – soil – 15
65. *Neckera complanata* (Hedw.) Huebener – tree base – 46
66. *Orthotrichum affine* Schrad. ex Brid. – tree bark – 16
67. *Orthotrichum anomalum* Hedw. – rock and concrete fence – 8, 17
68. *Orthotrichum cupulatum* Hoffm. ex Brid. – rock – 15, 27
69. *Orthotrichum lyellii* Hook. & Taylor – tree bark – 24
70. *Orthotrichum tenellum* Bruch ex Brid. – tree bark – 25
71. *Oxyrrhynchium hians* (Hedw.) Loeske – soil – 32
72. *Phascum cuspidatum* Hedw. – soil – 12, 15, 19, 20, 33, 43
73. *Plagiomnium undulatum* (Hedw.) T. J. Kop. – soil – 25
74. *Plagiothecium nemorale* (Mitt.) A. Jaeger – soil – 16
75. *Platyhypnidium riparioides* (Hedw.) Dixon – in and by the water – 1, 13, 25, 35
76. *Pleurochaete squarrosa* (Brid.) Lindb. – soil – 3, 8, 11, 16, 17, 21, 22, 24, 26, 32, 38, 43, 44
77. *Pogonatum aloides* (Hedw.) P. Beauv. – soil – 25, 26
78. *Pohlia melanodon* (Brid.) A. J. Shaw – soil by the water – 13
79. *Polytrichum juniperinum* Hedw. – soil – 25
80. *Polytrichum piliferum* Hedw. – soil – 26
81. (a) *Protobryum bryoides* (Dicks.) J. Guerra & M. J. Cano – soil – 15, 24
82. *Pseudocrossidium hornschuchianum* (Schultz) R. H. Zander – soil – 1, 2, 3, 5, 6, 12, 13, 18, 19, 22, 23, 27, 29, 35, 36, 38, 42, 46
83. *Pseudocrossidium revolutum* (Brid.) R. H. Zander – soil – 26, 33, 38, 46



84. *Pseudoscleropodium purum* (Hedw.) M. Fleisch – soil – 16
85. *Pterogonium gracile* (Hedw.) Sm. – rock – 25
86. (a) *Pterygoneurum ovatum* (Hedw.) Dixon – soil – 13
87. *Racomitrium canescens* (Hedw.) Brid. – siliceous rock – 25
88. *Rhynchostegiella curviseta* (Brid.) Limpr. – wet rocks – 26
89. + *Rhynchostegiella litorea* (De Not.) Limpr. – wet rocks – 25
90. *Rhynchostegiella tenella* (Dicks.) Limpr. – shaded rock crevices – 33, 34
91. *Schistidium apocarpum* (Hedw.) Bruch & Schimp. – rocks – 16, 24
92. + *Schistidium brunnescens* Limpr. – rock – 33
93. *Schistidium crassipilum* H. H. Blom – rock, concrete – 33, 37, 46
94. *Schistidium elegantulum* H. H. Blom – rock – 30
95. + *Schistidium helveticum* (Schkuhr) Deguchi – rock – 17, 34
96. *Scleropodium touretii* (Brid.) L. F. Koch – soil and slopes – 11, 12, 16, 23, 24, 25, 26, 27, 28, 29, 30, 31, 37, 45
97. + *Syntrichia calcicola* J. J. Amann – soil and rocks – 32, 36, 46
98. *Syntrichia princeps* (De Not.) Mitt. – soil – 40
99. *Syntrichia ruralis* (Hedw.) F. Weber & D. Mohr – 7, 8, 9, 14, 25, 29, 33, 40, 43
100. + *Syntrichia ruralis* var. *ruraliformis* (Besch.) Delogne – soil – 17, 21, 45
101. *Thamnobryum alopecurum* (Hedw.) Gangulee – tree base – 25
102. *Tortella flavovirens* (Bruch) Broth. – soil – 16
103. *Tortella nitida* (Lindb.) Broth. – rocks – 26
104. *Tortella tortuosa* (Hedw.) Limpr. – rocky slope – 35
105. (a) *Tortula inermis* (Brid.) Mont. – soil – 25
106. (a) *Tortula lanceola* R. H. Zander – soil – 18, 36, 37
107. *Tortula muralis* Hedw. – rocks, concrete – all
108. + *Tortula muralis* var. *aestiva* Hedw. – rock – 26
109. *Tortula subulata* Hedw. – soil – 25
110. + *Tortula schimperi* M. J. Cano, O. Werner & J. Guerra – soil – 10
111. *Trichostomum brachydontium* Bruch – soil – 6, 9, 11, 17, 29, 31, 32, 38, 40, 44
112. *Trichostomum crispulum* Bruch – 4, 5, 8, 13, 14, 23, 39
113. *Weissia brachycarpa* (Nees & Hornsch.) Jur. – soil – 16
114. *Weissia controversa* Hedw. – soil – 26
115. *Weissia longifolia* Mitt. – soil – 16
116. *Zygodon rupestris* Schimp. ex Lorentz – tree bark and base – 16

Among the 138 bryophytes recorded in the study area, one species (*Conocephalum salebrosum*) is newly recorded for Greece. Three taxa are new for the mainland of Greece: *Fissidens serrulatus* (previously reported only from Rhodos island), *Rhynchostegiella litorea* and *Schistidium helveticum* (both previously reported only from the island of Crete). *Didymodon umbrosus* represents here the second record for Greece, since it has been only reported from Sterea Hellas (Athens, around Acropolis; Blockeel *et al.*, 2002).





Eighteen taxa are new to the north-eastern floristic region of Greece (marked with a + in the list), and 23 are first reported from Chalkidiki (marked with an "a" or + in the list).

Some species common in other parts of the Mediterranean, but with limited reports in Greece and SE Europe were recorded: *Crossidium crassinerve*, previously reported from Crete (Düll, 1966), Peloponnisos (Blockeel, 1991), and Sterea Hellas (Unger, 1862; Haussknecht, 1899; Rungby, 1971); *Protobryum bryoides*, reported from the Ionian islands and Sterea Hellas (Hararas, 1976), and recently from another site in NE Greece (Strymon river delta; Papp *et al.*, 2002); *Pterygoneurum ovatum*, here first reported from Chalkidiki, and earlier reported from the Ionian islands (Mazziari, 1851), Sterea Hellas (Geheeb, 1886) and North East Greece (near Xanthi; Taylor, 1952); *Schistidium brunnescens*, confirmed only from Crete (Düll, 1966), although it was included but only as synonym in Preston (1984b); *Tortula muralis* var. *aestiva*, previously reported from different islands (Crete, the Aegean islands and Rhodos *in* Düll, 1995a, the Ionian islands *in* Hararas, 1976), and in the mainland only from Sterea Hellas (Schiffner & Baumgartner, 1919). *Schistidium elegantulum*, after its first and unique report for Greece from Chalkidiki (Cholomontas Mt area; Lüth, 2003) is again recorded in Chalkidiki, but by the sea area of Ouranoupolis. Finally *Weissia longifolia*, recorded in this study at Metamorfosi area (sea level) confirms its presence in Chalkidiki. It had been previously reported from the Ionian islands (Düll, 1995a; Damm *et al.* pers. comm.), and from NE Greece as cf. at Cholomontas Mt area/ Chalkidiki (Erzberger, 2006).

Some of the recorded taxa also deserve special attention due to their threat status and rareness.

Conocephalum conicum is a common and wide-spread species, and the newly described sister species *C. salebrosum* (Szweykowski *et al.*, 2005) is easy to overlook. Among the 13 recorded populations of *Conocephalum* in the study area, only one specimen belonged to *C. salebrosum*. In SE Europe it is known to occur in Romania and Bosnia-Herzegovina (Sabovljević & Natscheva, 2006).

Barbula commutata (syn. *B. sardoa* Schimp., *B. convoluta* var. *commutata*) is a well defined taxon (Frahm & Ahmed, 2004a, b), considered to be either a variety of *B. convoluta* or a separate species better adapted to drought (Frahm, 2000; Hill *et al.*, 2006). In the study area both species occur and are easily overlooked in dry state.

Campylopus oerstadianus is a relict circum-tethian species (Frahm, 1984) with only some 12 populations known in Europe, of which only one, including two subpopulations, was found in the study area and in SE Europe. Previous collections were from the area of Cholomontas Mt. area, Vourvourou (Düll, 1995b), and Neos Marmaras-Parthenon (Erzbeger, 2006). In this study a new locality in Chalkidiki area is added. Its status in Europe is rare (R) with insufficient knowledge on its population stages (ECCB, 1995). The species deserves a special conservation status in Greece as well.

Gymnostomum lanceolatum, has been described from Spain (Cano *et al.*, 1994) and later recorded in other Mediterranean countries: Italy (Aleffi *et al.*, 2004), Croatia (Sabovljević, 2006), Turkey (Kučera, 1998), and southern Greece: Peloponnisos, Sterea Hellas and East Central Greece (Blockeel *et al.*, 2002). Some authors consider it a variety of *G. calcareum* Nees & Hornsch. (Sérgio, 2006). Similarly, *Didymodon sicculus* appears to turn from a Spanish endemic (Cano *et al.*, 1996) to a Mediterranean species by its records in other Mediterranean countries: Montenegro (Cvetić & Sabovljević, 2004), Italy (Aleffi *et al.*, 2003a-b, Puglisi *et al.*, 2004), Turkey (Papp & Sabovljević, 2003). In Greece it was recorded



in Peloponnisos and Sterea Hellas (Blockeel *et al.*, 2002) and this is the second record for the country. It is likely to occur in other areas in Greece. Finally, *Tortula schimperi* recently separated from *T. subulata* complex (Cano *et al.*, 2005) is recorded as new for NE Greece.

Acknowledgement. The authors acknowledge two anonymous referees and to V. Mazimpaka (Madrid) for the comments on previous version of manuscript.

REFERENCES

- ALEFFI M., SABOVLJEVIĆ M. & TACCHI R., 2003a — *Didymodon sicculus* M. J. Cano, Ros, García-Zamora & J. Guerra (Pottiaceae, Musci), new to Italy. *Cryptogamie, Bryologie* 24 (1): 49-51.
- ALEFFI M., SABOVLJEVIĆ M. & TACCHI R., 2003b — Bryoflora of Monte Conero Regional Park (Marche, Italy). *Flora Mediterranea* 13: 151-171.
- ALEFFI M., SABOVLJEVIĆ M. & TACCHI R., 2004 — *Gymnostomum lanceolatum* M.J. Cano, Ros & J. Guerra (Pottiaceae, Musci), new to Italy. *Cryptogamie, Bryologie* 25 (2): 175-177.
- ATHANASIADIS N., 1977 — *Sphagnum contortum* Schultz, *Sph. subsecundum* Nees, *Sph. palustre* L. and *Sph. squarrosum* Pers (Cr.) four new species of the Greek flora. *Aristotle University of Thessaloniki. Scientific Annals of Agronomic and Forestry School* 10: 263-281 [in Greek, with German summary].
- BALAFOUTIS Ch., 1977 — Contribution to the study of the climate of Macedonia and West Thrace. Ph.D Thesis. Aristotle University of Thessaloniki. Thessaloniki, Greece [in Greek].
- BLOCKEEL T.L., 1991 — The bryophytes of Greece: new records and observations. *Journal of bryology* 16: 629-640.
- BLOCKEEL T.L., 2001 — Winter on a Greek island, the bryophytes of Evvia. *Bulletin of the British bryological society* 76: 26-28.
- BLOCKEEL T.L., ROS R.M., SABOVLJEVIĆ M., CANO M.J., GALLEGOS M.T. & MUÑOZ J., 2002 — New and interesting bryophyte records for Greece. *Cryptogamie, Bryologie* 23(2): 149-155.
- BLOCKEEL T.L. & RAMEL G., 2006 — New national and regional bryophyte records: *Physcomitrella patens* (Hedw.) Bruch & Schimp. *Bryological Notes. Journal of bryology* 28 (3): 273.
- CANO M.J., ROS R.M. & GUERRA J., 1994 — *Gymnostomum lanceolatum* sp. nov. (Pottiaceae, Musci) von der Iberischen Halbinsel. *Nova Hedwigia* 59: 143-146.
- CANO M.J., ROS R.M., GARCÍA-ZAMORA P. & GUERRA J., 1996 — *Didymodon sicculus* sp. nov. (Bryopsida, Pottiaceae) from the Iberian Peninsula. *The bryologist* 99: 401-406.
- CANO M.J., WERNER O. & GUERRA J., 2005 — A morphometric and molecular study in *Tortula subulata* complex (Pottiaceae, Bryophyta). *Botanical journal of the Linnean society* 149: 333-350.
- CARRATELLO A. & ALEFFI M., 1998 — *Gigaspernum mouretii* Corb. (Gigaspermaceae, Musci), a new species from Italy. *Acta botanica Malacitana* 23: 203-207.
- CVETIĆ T. & SABOVLJEVIĆ M., 2004 — New and interesting bryophyte records for Montenegro. *Phytologia balcanica* 10 (2-3): 67-69.
- DAFIS S., PAPASTERGIADOU E., GEORGHIOU K., BABALONAS D., GEORGIADIS T., PAPAGEORGIOU M., LAZARIDOU TH. & TSIAOUSSI V., 1996 — Directive 92/43/EEC The Greek "Habitat" Project NATURA 2000: An Overview. Life Contract B4 -3200/94/756, Commission of the European Communities DG XI, The Goulandris Natural History Museum, GreekBiotope/Wetland Centre, 917 p.
- DÜLL R., 1966 — Moosflora und -vegetation auf Kreta. *Revue bryologique et lichenologique* 34 (3-4): 732-761.
- DÜLL R., 1995a — Survey of the Bryophytes of Greece. *Bryologische Beiträge* 10: 1-125.
- DÜLL R., 1995b — Bryophytes of the Chalkidiki (incl. Sithonia and Athos) and the Rhodopi mountains. *Bryologische Beiträge* 10: 126-142.
- DÜLL R., 1996 — The current state of bryophyte investigation in the Mediterranean area. *Bocconeia* 5: 271-278.
- ECCB, 1995 — *Red Data Book of European Bryophytes*. Trondheim, The European Committee for Conservation of Bryophytes, 291 p.



- ERZBERGER P. 2006 — Contribution to the bryophyte flora of Chalkidiki, Greece. *Willdenowia* 36: 515-525.
- FRAHM J.-P. 1984 — Phytogeography of European *Campylopus* Species. *Proceedings of the 3rd Meeting of Bryologists from Central and East Europe*, Prague, pp. 191-212.
- FRAHM J.-P., 2000 — *Barbula convoluta* var. *commutata*, eine verkannte Sippe. *Bryologische Rundbriefe* 33: 7.
- FRAHM J.-P. & AHMED J., 2004a — *Barbula sardoa* (Schimp.) J.-P. Frahm, a new name for *Barbula convoluta* Hedw. var. *commutata* (Jur.) Husn. *Journal of bryology* 26: 29-35.
- FRAHM J.-P. & AHMED J., 2004b — *Barbula sardoa* (Schimp.) J.-P. Frahm, a new name for *Barbula convoluta* Hedw. var. *commutata* (Jur.) Husn. [Erratum]. *Journal of bryology* 26: 244.
- GALLEGU M.T. & CANO M.J., 1998 — *Aloina brevirostris* (Hook. & Grev.) Kindb., new for Greece, the Iberian Peninsula and the Canary Islands. Bryological Notes. *Journal of bryology* 20: 245-246.
- GAMISANS J. & HÉBRARD J.P., 1980 — A propos de la végétation des forêts en Grèce du nord-est (Macédoine orientale et Thrace occidentale). *Documents phytosociologiques* n.s., 5: 243-289.
- GANIATAS K.A., 1938 — Contribution to the knowledge of Bryophytes of Macedonia. *Annal. Scient. University of Thessaloniki, School of Natural and Mathematical Sciences*, 4: 75-93 [in Greek].
- GEHEEB A., 1886 — Bryologische Fregmente. III. C. Griechische Laubmoose. *Flora* 69: 324-323.
- GEISSLER P., 1977 — Zur Moos- und Flechtenflora Nordgriechenlands. *Bauhinia* 6(1): 189-213.
- GIANNITSAROS A. & KOUMPLI-SOVANTZI L., 1990 — *Sphagnum cuspidatum* Ehrh. ex Hoffm., a new species of the Hellenic flora. *Proceedings of the 12th Scientific Congress of the Hellenic Society of Biological Sciences, Mytilini*, pp. 41-44 [in Greek, with English summary].
- HARARAS K., 1976 — *Ereynai epi ton bryophyton tis nisoy Kerkyras* (*Study of bryophytes at the Corfu island. Chorological, ecological, phytogeographical study*). PhD thesis. School of Natural Sciences, University of Athens, 160 p. (+3 maps) [in Greek].
- HAUSSKNECHT H.K., 1899 — Symbolae ad floram graecam. Aufzählung der im Sommer 1885 in Griechenland gesammelten Pflanzen. *Mitteilungen thüringische botanische Vereins*, Neue Folge, 13: 18-77.
- HEDDERSON T.A. & BLOCKEEL T.L., 2006 — *Oncophorus dendrophilus*, a new moss species from Cyprus and Crete. *Journal of bryology* 28(4): 357-359.
- HILL M.O., BELL N., BRUGGEMANN-NANNENGA M.A., BRUGUES M., CANO M.J., ENROTH J., FLATBERG K.I., FRAHM J.-P., GALLEGU M.T., GARILLETTI R., GUERRA J., HEDENÄS L., HOLYOAK D.T., HYVÖNEN J., IGNATOV M.S., LARA F., MAZIMPAKA V., MUÑOZ J. & SÖDERSTRÖM L., 2006 — An annotated checklist of the mosses of Europe and Macaronesia. *Journal of bryology* 28(3): 198-267.
- KUČERA J., 1998 — *Gymnostomum lanceolatum* M. J. Cano, Ros & J. Guerra, aslo in Turkey and Croatia. *Journal of bryology* 20: 515-516.
- LARA F., BLOCKEEL T.L., GARILLETTI R. & MAZIMPAKA V., 2003 — Some interesting *Orthotrichum* species from mainland Greece and Evvia. *Journal of bryology* 25: 129-134.
- LÜTH M., 2002 — *Cinclidous confertus* (*Musci, Cinclidotaceae*), a new species from Greece. *Cryptogamie, Bryologie* 23(1): 11-16.
- LÜTH M., 2003 — Moose in der Umgebung des Dorfes Vikos, im Vikos-Aoos Nationalpark (NW Griechenland). *Archive for bryology* 2: 1-25.
- MAIER E., 2002 — *Grimmia dissimulata* E. Maier sp. nova, and the taxonomic position of *Grimmia trichophylla* var. *meridionalis* Müll. Hal. (*Musci, Grimmiaceae*). *Candollea* 56(2): 281-300.
- MAVRONMATIS G., 1972 — Description of a turf-type vegetation in Elatia (Kara Dere) Drama. *Dasos* (55-56): 26-27 [in Greek, with English summary].
- MAZZIARI A.D., 1851 — Flora Septinsularis. Published only as extracts in STEFANIDES, Th.(1948). A survey of the freshwater biology of Corfu and of certain other regions of Greece. *Praktika Hellenic Hydrobiology Institute, Academy of Science, Athens*, 2: 1-263, Appendix 1, pp. 187-201 [in Greek].
- MOUNTRAKIS D., 1985 — *The Geology of Greece*. Aristotle University of Thessaloniki. Thessaloniki, Greece, University Studio Press [in Greek].
- PAPP B. & BABALONAS D., 1996 — Bryophyte flora of some maquis and phrygana stands in Greece. *Bios (Macedonia, Greece)* 4: 115-122.
- PAPP B., 1998 — Investigation of the bryoflora of some streams in Greece. *Studia botanica hungarica* 29: 59-67.
- PAPP B., LÖKÖS L., RAJCZY M., CHATZINIKOLAKI E. & DAMANAKIS M., 1998a — Bryophytes and lichens of some phrygana and maquis stands of Crete (Greece). *Studia botanica hungarica* 29: 69-78.



- PAPP B., TSAKIRI E. & BABALONAS D., 1998b — Bryophytes and their environmental conditions at Enipeas (Mt Olympos) and Lykorrema (Mt Ossa) streams (Greece). In: Tsekos I. & Moustakas M. (eds), 'Progress in Botanical Research'. Proc. 1st Balkan Botanical Congress, Thessaloniki, Greece. Dordrecht, etc., pp. 129-132.
- PAPP B., 2002 — New records of bryophytes from a saline area of Greece. *Studia botanica hungarica* 33: 21-24.
- PAPP B., TSAKIRI E. & KARAGIANNAKIDOU V., 2002 — Bryophytes of two alkali areas of Northern Greece. *Proceedings of the 9th Scientific Congress of the Hellenic Botanical Society*, Kefalonia, Argostoli, pp. 233-240.
- PAPP B., 2003 — New records of bryophytes from Crete. *Studia botanica hungarica* 34: 27-31.
- PAPP B. & SABOVLJEVIĆ M., 2003 — Contribution to the bryophyte flora of Turkish Thrace. *Studia botanica hungarica* 34: 43-54.
- PARENT G.H. & DE ZUTTERE P., 2006 — *Sphagnum teres* (Schimp.) Ångstr., nouveau pour la Grèce, dans une pozzine de la Macédoine septentrionale grecque. *Newellia* 30: 2-8.
- PODPĚRA J., 1922 — Ad Bryophytorum Haemi peninsulae cognitionem additameuntum. *Acta botanica bohemica* I: 5-25.
- POLITIS J., 1953 — Contribution a l'étude de la flore de la Chalkidique. *Bryophyta* pp: 49-53. Pragmateiai Academeias Atheneion, Academy of Athens, 19: 1-97.
- PRESTON C.D., 1981 — A check-list of Greek liverworts. *Journal of bryology* 11: 537-553.
- PRESTON C.D., 1984a — A check-list of Greek liverworts: Addendum. *Journal of bryology* 13: 97-100.
- PRESTON C.D., 1984b — A check-list of Greek mosses. *Journal of bryology* 13: 43-95.
- PUGLISI M., PRIVITERA M. & PREZZAVENTO A.D., 2004 — More about the occurrence of *Didymodon sicculus* Cano, Ros, Garcia-Zamora & Guerra (Musci, Pottiaceae) in Italy. *Braun-Blanquetia* 34: 47-49.
- REIMERS H., 1957 — Einige bemerkenswerte Mosse des östlichen Mediterrangebiets. *Willdenowia* 1(5): 689-703.
- ROS R.M., MAZIMPANA V., ABOU-SALAMA U., ALEFFI M., BLOCKEEL T.L., BRUGUÉS M., CANO M.J., CROS R.M., DIA M.G., DIRKSE G.M., EL SAADAWI W., ERDAĞ A., GANEVA A., GONZÁLEZ-MANCEBO J.M., HERRNSTADT I., KHALIL K., KÜRSCHNER H., LANFRANCO E., LOSADA-LIMA A., REFAI M.S., RODRÍGUEZ-NUÑEZ S., SABOVLJEVIĆ M., SÉRGIO C., SHABBARA H., SIM-SIM M., SÖDERSTRÖM L. 2007 — Hepatics and Anthocerotes of the Mediterranean, an annotated checklist. *Cryptogamie, Bryologie* 28: 351-437.
- RUNGBY S. 1971 — A contribution to the bryophytic flora of Attica and Peloponnese. *Botaniske notiser* 124: 228-230.
- SABOVLJEVIĆ M., 2004 — Comparison of the bryophyte flora of the three southern European mainlands: the Iberian, the Apennine and the Balkan peninsulas. *Braun-Blanquetia* 34: 21-28.
- SABOVLJEVIĆ M., 2006 — The check list of mosses of Croatia. *Archive of biological sciences* 58: 45-53.
- SABOVLJEVIĆ M. & NATCHEVA R., 2006 — Check list of the liverworts and hornworts of South-Eastern Europe. *Phytologia balcanica* 12(2): 169-180.
- SABOVLJEVIĆ M., GANEVA A., TSAKIRI E. & STEFANUT S., 2001 — Bryology and bryophyte protection in the south-eastern Europe. *Biological conservation* 101: 73-84.
- SAUER E., 1995 — Beitrag zur Moosflora Griechenlands. *Bryologische Beiträge* 10: 209-217.
- SCHIFFNER V. & BAUMGARTNER J., 1919 — Beiträge zur Kenntnis der Flora Griechenlands. B. Leber- und Laubmoose. *Verhandlungen der Zoologisch-Botanischen Gesellschaft Wien* 69: 313-341.
- SÉRGIO C., 2006 — A review of the *Gymnostomum calcareum* Nees & Hornsch. complex (Bryopsida: Pottiaceae) in southern Europe and the Macaronesian Islands, including *G. calcareum* var. *atlanticum* var. nov. *Journal of bryology* 28 (1): 38-45.
- SIBTHORP J. & SMITH J.E., 1806-1813 — *Florae Graecae Prodromus*. London.
- SÖDERSTRÖM L., HALLINGBÄCK T., HODGETTS N., RAEYMAEKERS G., SCHUMACKER R., SÉRGIO C., STEWART N. & VÁŇA J., 1998 — State of knowledge of the bryoflora of Europe as illustrated by the hepatic flora. *Lindbergia* 23: 28-32.
- SZWEYKOWSKI J., BUCZKOWSKA K. & ODRZYKOSKI I. J., 2005 — *Conocephalum salebrosum* (Marchantiopsida, Conocephalaceae) - a new Holarctic liverwort species. *Plant systematic and evolution* 253: 133-158.
- TAYLOR J., 1952 — A contribution to the bryophytic flora of the Nearer East. *Kew bulletin* 1952: 45-60.
- TSAKIRI E. & BABALONAS D., 2002 — Bryophyte flora of the drainage system of Grevena Prefecture (West Macedonia, Greece). *Proceedings of the 9th Scientific Congress of the Hellenic Botanical Society*, Cephallonia, Greece, pp. 319-324 [in Greek, with English abstract].
- TSAKIRI E., KARAGIANNAKIDOU V. & BABALONAS D. †, 2005 — Contribution to the study of the bryophytic flora of Northern Greece (Bryophyte flora of the hydrographic network



of Ano Aliakmonas, West Macedonia). pp. 7. *Proceedings of the 10th Scientific Congress of the Hellenic Botanical Society*, Ioannina, Greece. (in press) [in Greek, with English abstract].

TSAKIRI E., PAPP B. & SZURDOKI E., 2006 — New records on the bryophytes of Mt Voras, North Greece. *Studia botanica hungarica* 37: 145-155.

TURLAND N.J. & WILSON C.C., 1995 — *Sphagnum auriculatum* Schimp.: a genus new to the bryoflora of Crete. *Journal of bryology* 18: 827-828.

UNGER F., 1862 — Wissenschaftliche Ergebnisse einer Reise in Griechenland und in der Ionischen Inseln. Wien, pp. 108-115.

VAJDA L., 1959 — Moose aus Griechenland. *Annales historico-naturales musei nationalis Hungarici* 51: 197-199.

ZOLLER H., GEISSLER P. & ATHANASIADIS N., 1977 — Beiträge zur Kenntnis der Wälder, Moos - und Flechtenassoziationen in den Gebirgen Nordgriechenlands. *Bauhinia* 6(1): 215-255.

P R O O O F