

Length-Weight Relationships of Freshwater Fishes in Greece

P.K. Kleanthidis, A.I. Sinis and K.I. Stergiou

Abstract

Length-weight relationships were calculated for nine fish species from Lake Volvi (Macedonia, Hellas), caught with gillnets of five different mesh sizes between October 1995 and October 1996. In addition, length-weight relationships for 24 Greek freshwater fish species and one hybrid were also obtained from the literature. The values of the exponent of the length-weight relationships for all fish species examined ranged between 2.14 and 3.70 (mean = 3.12; SE = 0.032), and the median value was 3.19.

Introduction

In fisheries research, length-weight relationships are important for the estimation of weight where only length data are available and as an index of the condition of the fish (Pauly 1993; Petrakis and Stergiou 1995; Goncalves et al. 1997). In this study, the parameters of the length-weight relationships for nine fish species collected from Lake Volvi are estimated. In addition, the parameters of the length-weight relationships for 24 Greek freshwater fish species and one hybrid collected from the literature are also presented.

Materials and Methods

Samples were collected monthly from October 1995 to October 1996 from Lake Volvi (Macedonia, Hellas), using gillnets of mesh sizes 14, 18, 22, 26 and 30 mm (nominal bar length). Each net was 100 m long and 2 m deep. All fish were preserved in 10% formalin solution immediately after capture. In the laboratory, the total length was measured to the nearest mm and the total weight was weighed to the nearest g.

The parameters *a* and *b* of the length-weight relationship ($W=aL^b$)

were calculated for each species (after log-transformation). All weights are expressed in g and all lengths in cm.

Results and Discussion

A total of nine species were caught in Lake Volvi, namely: *Abramis brama*, *Alburnus alburnus*, *Alosa macedonica*, *Carassius auratus gibelio*, *Chalcalburnus chalcoides*, *Cyprinus carpio*, *Rutilus rutilus*, *Scardinius erythrophthalmus* and *Vimba melanops*. The parameters *a* and *b* of the length-weight relationships are shown in Table 1. The sample size ranged from 11 individuals for *C. carpio* to 4825 for *A. macedonica*. The values of *b* ranged from 2.67 for *A. alburnus* to 3.48 for *S. erythrophthalmus*.

The values of the parameters of the length-weight relationships of the 24 Greek freshwater fish species and one hybrid (from ten lakes, seven rivers and two lagoons), collected from the literature (a total of 85 length-weight relationships) are shown in Table 2.

Overall (Tables 1 and 2), the value of *b* ranged from 2.14 for *Leuciscus 'svallize'* of Lake Kremasta, to 3.70 for female *A. alburnus* of Lake Mikri Prespa. The

Table 1. Parameters of the relationships ($W=aL^b$) between total weight (in g) and total length (in cm) of nine fish species, collected monthly from October 1995 to October 1996, from Lake Volvi. *N*, sample size; *min* and *max*, minimum and maximum total lengths in cm, respectively; *SE(b)*, standard error of *b*; *r*, correlation coefficient.

Species ¹	N	Length		a	b	SE(b)	r
		min	max				
<i>Abramis brama</i>	449	10.7	30.5	0.0111	2.97	0.036	0.97
<i>Alburnus alburnus</i>	86	9.4	15.6	0.0185	2.70	0.127	0.92
<i>Alosa macedonica</i>	4825	8.3	21.4	0.0176	2.71	0.011	0.96
<i>Carassius auratus gibelio</i>	102	8.2	25.2	0.0142	3.11	0.057	0.98
<i>Chalcalburnus chalcoides</i>	67	12.9	21.5	0.0029	3.41	0.076	0.98
<i>Cyprinus carpio</i>	11	7.8	18.1	0.0383	2.67	0.090	0.99
<i>Rutilus rutilus</i>	4338	9.3	26.0	0.0074	3.14	0.013	0.96
<i>Scardinius erythrophthalmus</i>	75	10.6	21.1	0.0036	3.48	0.057	0.99
<i>Vimba melanops</i>	22	14.5	22.8	0.0043	3.33	0.132	0.98

¹ Species are listed alphabetically. Species names according to Economidis and Sinis (1982) and Economidis (1991).

Table 2. Parameters of the relationships ($W=aL^b$) between weight (in g) and length (in cm) of 24 Greek freshwater fish species and one hybrid, from ten lakes, seven rivers and two lagoons, collected from the literature. F, frequency of sampling (M, monthly; B, bimonthly; S, seasonal; O, one single sampling; U, undefined); sex (M, male; F, female; I, immature fish; C, sexes combined); weight (total, TW; net, NW); length (fork, FL; standard, SL; total, TL); N, sample size; length characteristics (min and max, minimum and maximum lengths in cm, respectively); SE(b), standard error of b; r, correlation coefficient.

Species	F	Sex	Weight	Length	N	Length characteristics		a	b	SE(b)	r	Area/Sampling period	Reference
						min	max						
<i>Abramis brama</i>	M	M	TW	FL	655	11.7	32.0	0.0082	3.18	-	-	Lake Volvi /Jan-Dec 1990	Valoukas et al. (1996)
<i>Abramis brama</i>	M	F	TW	FL	654	11.9	36.3	0.0075	3.21	-	-	Lake Volvi /Jan-Dec 1990	Valoukas et al. (1996)
<i>Alburnus alburnus</i>	M	M	TW	TL	1403	-	-	0.0049	3.23	0.035	0.93	Lake Koronia /Apr 1986-Sep 1988	Politou (1993)
<i>Alburnus alburnus</i>	M	F	TW	TL	1522	-	-	0.0029	3.43	0.027	0.95	Lake Koronia /Apr 1986-Sep 1988	Politou (1993)
<i>Alburnus alburnus</i>	M	C	TW	TL	2925	7.7	15.4	0.0038	3.33	0.019	0.95	Lake Koronia /Apr 1986-Sep 1988	Politou (1993)
<i>Alburnus alburnus</i>	U	M	-	FL	78	7.5	17.5	0.0021	3.64	-	0.97	Lake Mikri Prespa /Mar-Jun 1985	Crivelli and Dupont (1987)
<i>Alburnus alburnus</i>	U	F	-	FL	200	7.5	17.5	0.0018	3.70	-	0.98	Lake Mikri Prespa /Mar-Jun 1985	Crivelli and Dupont (1987)
<i>Alosa macedonica</i>	O	M	NW	TL	123	10.1	18.0	0.0179	2.66	0.056	0.97	Lake Volvi /Jun 1978	Sinis (1981)
<i>Alosa macedonica</i>	O	F	NW	TL	212	11.1	23.0	0.0168	2.69	0.037	0.98	Lake Volvi /Jun 1978	Sinis (1981)
<i>Alosa macedonica</i>	O	C	NW	TL	335	10.1	23.0	0.0157	2.72	0.024	0.98	Lake Volvi /Jun 1978	Sinis (1981)
<i>Aphanius fasciatus</i>	M	M	NW	TL	418	2.2	6.8	0.0094	3.31	0.035	0.97	Messolonghi lag. (Rebakia) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	F	NW	TL	493	2.0	7.0	0.0088	3.45	0.015	0.99	Messolonghi lag. (Rebakia) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	C	NW	TL	911	2.0	7.0	0.0088	3.43	0.015	0.99	Messolonghi lag. (Rebakia) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	M	NW	TL	218	2.0	5.0	0.0097	3.22	0.030	0.99	Eliotikon lag. (Astrovitsa) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	F	NW	TL	233	2.6	7.2	0.0087	3.45	0.040	0.99	Eliotikon lag. (Astrovitsa) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	C	NW	TL	451	2.0	7.2	0.0086	3.45	0.015	0.99	Eliotikon lag. (Astrovitsa) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	M	NW	TL	293	2.0	5.8	0.0100	3.09	0.040	0.97	Messolonghi lag. (Alykes) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	F	NW	TL	310	2.2	6.4	0.0100	3.16	0.035	0.98	Messolonghi lag. (Alykes) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Aphanius fasciatus</i>	M	C	NW	TL	648	2.0	6.4	0.0100	3.18	0.025	0.98	Messolonghi lag. (Alykes) /Apr 1989-Jan 1991	Leonardos (1996)
<i>Atherina boyeri</i>	U	C	TW	TL	2543	-	-	0.0049	3.21	-	0.98	Lake Trichonis /1988-90	Stourboudi et al. (1997)
<i>Barbus albanicus</i>	S	M	-	FL	149	5.0	20.1	0.0169	2.94	-	0.98	Lake Kremasta /Feb-Nov 1982	Daoulas and Economidis (1989)
<i>Barbus albanicus</i>	S	F	-	FL	175	5.0	28.8	0.0178	2.95	-	0.98	Lake Kremasta /Feb-Nov 1982	Daoulas and Economidis (1989)
<i>Barbus cyclolepis</i>	B	M	-	FL	-	5.0	26.0	0.0583	2.43	-	0.94	Rihios Stream /Feb-Dec 1984	Neophitou (1987)
<i>Barbus cyclolepis</i>	B	F	-	FL	-	5.0	26.0	0.0080	3.16	-	0.98	Rihios Stream /Feb-Dec 1984	Neophitou (1987)
<i>Barbus prespensis</i>	M	C	-	FL	144	8.0	30.0	0.0082**	3.20	-	0.99	Lake Mikri Prespa /Apr-Jun 1984, 1985, 1990-1994	Crivelli et al. (1996)
<i>Chalcaburnus belvica</i>	M	M	NW	FL	183	10.2	13.6	0.0058	3.23	0.070	0.96	Lake Mikri Prespa /Jul 1990-Jun 1991	Sinis and Petridis (1995)
<i>Chalcaburnus belvica</i>	M	F	NW	FL	186	10.6	22.0	0.0046	3.30	0.035	0.99	Lake Mikri Prespa /Jul 1990-Jun 1991	Sinis and Petridis (1995)
<i>Chalcaburnus belvica</i>	M	C	NW	FL	369	10.2	22.0	0.0067	3.18	0.030	0.98	Lake Mikri Prespa /Jul 1990-Jun 1991	Sinis and Petridis (1995)
<i>Chalcaburnus chalcoides</i>	M	M	NW	FL	90	9.1	20.0	0.0037	3.40	-	0.98	Lake Volvi /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Chalcaburnus chalcoides</i>	M	F	NW	FL	280	11.1	26.0	0.0035	3.41	-	0.98	Lake Volvi /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Chalcaburnus chalcoides</i>	M	C	NW	FL	370	9.1	26.0	0.0039	3.38	-	0.98	Lake Volvi /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Chalcaburnus chalcoides</i>	M	M	NW	FL	260	9.1	20.0	0.0026	3.56	-	0.99	Lake Vistonis /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Chalcaburnus chalcoides</i>	M	F	NW	FL	253	9.1	24.0	0.0031	3.49	-	0.99	Lake Vistonis /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Chalcaburnus chalcoides</i>	M	C	NW	FL	513	9.1	24.0	0.0029	3.52	-	0.99	Lake Vistonis /Sep 1983-Oct 1984	Kokkinakis (1992)
<i>Cyprinus carpio</i>	U	M	-	TL	21	-	-	0.0474**	2.65	-	-	Lake Vistonis /Sep-Dec 1973	Tsimenidis (1976)
<i>Cyprinus carpio</i>	U	F	-	TL	34	-	-	0.0140**	2.99	-	-	Lake Vistonis /Sep-Dec 1973	Tsimenidis (1976)
<i>Coregonus lavaretus</i>	O	M	NW	FL	58	24.3	33.8	0.0174	2.86	0.175	0.91	Lake Tavropos /Nov 1987	Sinis and Petridis (1993)
<i>Coregonus lavaretus</i>	O	F	NW	FL	66	24.3	33.8	0.1203	2.27	0.140	0.89	Lake Tavropos /Nov 1987	Sinis and Petridis (1993)
<i>Coregonus lavaretus</i>	O	M	NW	FL	69	24.3	34.8	0.0126	2.95	0.135	0.94	Lake Vegeritis /Nov 1987	Sinis and Petridis (1993)
<i>Coregonus lavaretus</i>	O	F	NW	FL	58	24.3	34.8	0.0097	3.02	0.160	0.93	Lake Vegeritis /Nov 1987	Sinis and Petridis (1993)
<i>Knipowitschia caucasica</i>	M	M	NW	TL	230	-	2.91	0.0040	3.25	0.008	0.86	Evos River (Delta) /Feb 1983-Feb 1984	Kevrekidis et al. (1990)

Species	F	Sex	Weight	Length	N	Length characteristics		a	b	SE(b)	r	Area/Sampling period	Reference
						min	max						
<i>Knipowitschia caucasica</i>	M	F	NW	TL	158	-	4.11	0.0034	3.43	0.009	0.93	Evros River (Delta) / Feb 1983-Feb 1984	Kevrekidis et al. (1990)
<i>Knipowitschia caucasica</i>	M	I	NW	TL	9	1.28	1.70	0.0031	2.79	0.409	0.67	Evros River (Delta) / Feb 1983-Feb 1984	Kevrekidis et al. (1990)
<i>Knipowitschia caucasica</i>	M	C	NW	TL	397	1.28	4.11	0.0034	3.46	0.004	0.91	Evros River (Delta) / Feb 1983-Feb 1984	Kevrekidis et al. (1990)
<i>Ladigesocypris ghigii</i>	M	M	TW	-	133	-	-	0.0109	3.22	-	0.98	Loutani stream (Rhodes)/Feb 1991-Jan 1992	Corsini and Karantoni (1993)
<i>Ladigesocypris ghigii</i>	M	F	TW	-	132	-	-	0.0116	3.19	-	0.99	Loutani stream (Rhodes)/Feb 1991-Jan 1992	Corsini and Karantoni (1993)
<i>Ladigesocypris ghigii</i>	M	M	TW	-	146	-	-	0.0128	3.14	-	0.99	Gadoura stream (Rhodes)/Feb 1991-Jan 1992	Corsini and Karantoni (1993)
<i>Ladigesocypris ghigii</i>	M	F	TW	-	94	-	-	0.0137	3.12	-	0.99	Gadoura stream (Rhodes)/Feb 1991-Jan 1992	Corsini and Karantoni (1993)
<i>Leuciscus cephalus</i>	U	C	-	FL	489	8.9	30.5	0.0027	3.44	-	0.99	Rihios Stream /Apr-Jun 1984	Neophitou (1993b)
<i>Leuciscus cephalus</i>	M	C	TW	SL	220	0.92	6.01	0.0123	3.23	-	0.99	Myrtia Stream /May 1989-May 1990	Economou et al. (1991a)
<i>Leuciscus 'svalize'</i>	O	M	TW	FL	36	7.0	20.0	0.0946	2.29	-	0.97	Lake Kremasta /Feb 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	F	TW	FL	88	7.0	20.0	0.0280	2.72	-	0.95	Lake Kremasta /Feb 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	M	TW	FL	145	7.0	21.0	0.0419	2.55	-	0.94	Lake Kremasta /Apr 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	F	TW	FL	146	7.0	21.0	0.0067	3.23	-	0.96	Lake Kremasta /Apr 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	M	TW	FL	111	4.0	21.0	0.0117	3.04	-	0.99	Lake Kremasta /Jun 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	F	TW	FL	97	4.0	21.0	0.0106	3.08	-	0.99	Lake Kremasta /Jun 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	M	TW	FL	111	6.0	19.0	0.0185	2.86	-	0.99	Lake Kremasta /Aug 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	F	TW	FL	130	6.0	19.0	0.0150	2.93	-	0.99	Lake Kremasta /Aug 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	M	TW	FL	47	7.0	22.0	0.0125	3.01	-	0.99	Lake Kremasta /Nov 1982	Economou et al. (1991b)
<i>Leuciscus 'svalize'</i>	O	F	TW	FL	137	7.0	22.0	0.0137	2.14	-	0.96	Lake Kremasta /Nov 1982	Economou et al. (1991b)
<i>Perca fluviatilis</i>	U	C	TW	TL	317	13.0	22.9	0.0229	2.83	-	0.93	Lake Doirani /1989-1992	Neophitou (1993b)
<i>Perca fluviatilis</i>	U	C	NW	SL	709	6.2	20.3	0.0127	3.17	-	-	Lake Koronia /1975-1976	Papageorgiou (1977)
<i>Pseudorasbora parva</i>	U	C	-	FL	245	6.1	9.5	0.0078	3.27	-	0.99	Lake Mikri Prespa /Apr-Jun 1984, Apr-Jun 1985, 1990-1992	Roseochi et al. (1993)
<i>Phoxinellus pleurobipunctatus</i>	B	M	TW	FL	450	5.5	18.0	0.0110	3.08	-	-	Lake Kremasta /Feb-Nov 1982	Daoulas et al. (1987)
<i>Phoxinellus pleurobipunctatus</i>	B	F	TW	FL	598	4.0	22.0	0.0121	3.04	-	-	Lake Kremasta /Feb-Nov 1982	Daoulas et al. (1987)
<i>Rutilus rubilio</i>	M	M	TW	FL	161	8.0	18.0	0.0060	3.30	0.006	0.98	Lake Trichonis /Mar-May 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	F	TW	FL	246	8.0	26.0	0.0200	3.00	0.003	0.98	Lake Trichonis /Mar-May 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	M	TW	FL	163	9.0	20.0	0.0100	3.30	0.003	0.99	Lake Trichonis /Jun-Aug 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	F	TW	FL	208	8.0	23.0	0.0060	3.30	0.002	0.99	Lake Trichonis /Jun-Aug 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	M	TW	FL	138	9.0	18.0	0.0095	3.20	0.006	0.98	Lake Trichonis /Sep-Nov 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	F	TW	FL	144	5.0	20.0	0.0080	3.30	0.004	0.98	Lake Trichonis /Sep-Nov 1978	Daoulas (1981)
<i>Rutilus rubilio</i>	M	M	TW	FL	122	11.0	18.0	0.0111	3.20	0.008	0.98	Lake Trichonis /Dec-Feb 1978-1979	Daoulas (1981)
<i>Rutilus rubilio</i>	M	F	TW	FL	147	11.0	20.0	0.0079	3.20	0.004	0.98	Lake Trichonis /Dec-Feb 1978-1979	Daoulas (1981)
<i>Rutilus rubilio</i>	M	C	TW	FL	768	12.0	21.0	19.364?	3.14	-	0.93	Lake Pamvotis /Nov 1983-Dec 1984	Daoulas (1981)
<i>Rutilus rubilio</i>	U	C	-	FL	501	6.0	20.0	0.0074	3.32	-	0.99	Lake Mikri Prespa /Mar-Jun 1985	Neophitou et al. (1989)
<i>R. rubilio x A. alburnus</i>	U	M	-	FL	43	10.6	18.5	0.0100	3.11	-	0.97	Lake Mikri Prespa /Mar-Jun 1985	Crivelli and Dupont (1987)
<i>R. rubilio x A. alburnus</i>	U	F	-	FL	87	10.6	18.5	0.0054	3.33	-	0.97	Lake Mikri Prespa /Mar-Jun 1985	Crivelli and Dupont (1987)
<i>Rutilus rutilus</i>	O	M	TW	TL	97	7.2	23.0	0.0356	3.40	-	0.99	Lake Mikri Prespa /Mar-Jun 1985	Crivelli and Dupont (1987)
<i>Rutilus rutilus</i>	O	M	TW	TL	136	7.2	23.0	0.0215	3.61	-	0.99	Lake Volvi /Mar 1978	Papageorgiou (1979)
<i>Rutilus rutilus</i>	O	F	TW	TL	409	2.0	7.0	0.0068**	3.33	-	0.99	Lake Trichonis /Sep 1988-Apr 1991	Papageorgiou (1979)
<i>Salapia fluviatilis</i>	M	M	TW	TL	441	2.0	7.0	0.0102**	3.08	-	0.97	Lake Trichonis /Sep 1988-Apr 1991	Psarras et al. (1997)
<i>Salmo fluviatilis</i>	M	C	-	SL	1074	4.1	24.0	0.0041**	2.95	-	-	Aspropolamos Stream /Mar 1981-Feb 1982	Papageorgiou et al. (1983)
<i>Salmo trutta fario</i>	M	M	TW	TL	180	5.0	31.0	0.0194	2.86	-	0.98	Acheleos River /1978-1981	Klossa-Killa (1990)
<i>Salmo trutta</i>	M	F	TW	TL	128	9.0	29.0	0.0214	2.82	-	0.95	Acheleos River /1978-1981	Klossa-Killa (1990)
<i>Salmo trutta</i>	M	C	NW	FL	763	-	29.0	0.0295	2.80	-	0.98	Lake Pamvotis /Mar-Oct 1988	Neophitou (1993b)

Species are listed alphabetically. * Type of logarithm not specified, most probably base 10. ** Type of logarithm not specified, most probably natural.




Fig. 1. Box-Whisker plot of the exponent b of the length-weight relationships ($W=aL^b$) for all species examined (for values see Tables 1 and 2). The small open square inside the box represents the median value, the vertical line represents the minimum and the maximum values and the central box covers 50% of b values.

mean value of b for all species was 3.119 (SE = 0.032), the median value of b was 3.185 and 50% of the b values ranged between 2.950 and 3.325 (Fig. 1).

Some aspects are worthy of mention for Lake Volvi (Tables 1 and 2). The value of b did not change over time for *A. macedonica* (2.72 in 1978 and 2.71 in 1995-1996) and *C. chalcoides* (3.38 in 1983-1984 and 3.41 in 1995-1996). However, in the case of *A. macedonica* the estimate for 1978 was based on net weight and one sampling event whereas that for 1995-1996 was based on total weight and monthly sampling. Similarly, for *C. chalcoides*, the estimate in 1983-1984 was based on net weight and fork length while that for 1995-1996 was based on total length and total weight.

In contrast, the values of b for *A. brama* and *R. rutilus* seems to have changed considerably. For *A. brama*, the value of b decreased from 3.18 to 3.21 (depending on sex) in 1990 to 2.97 (for the sexes combined) in 1995-1996. This may be attributed to the use of fork length in the former period as opposed to the use of total length in

latter one (Tables 1 and 2). For *R. rutilus* the value of b also decreased from 3.40 to 3.61 (depending on sex) in 1978 (one sampling event) to 3.14 (sexes combined, monthly sampling) in 1995-1996.

Finally, the value of b for *A. alburnus* is considerably lower for Lake Volvi than for the nearby Lake Koronia (2.70 and 3.33, respectively).

Such changes in the values of b may be attributed either to differences in methodology, as described previously, or to factors like overfishing, food competition and trophic potential of the lakes, and require further study.

Acknowledgement

The authors wish to extend their gratitude to the fisher, Mr. T. Samaras, for assisting with the sampling.

References

Corsini, M. and A. Karantoni. 1993. Biological observations on two populations of *Ladigesocypris ghigii* (Pisces: Cyprinidae) in Rhodes, p. 295-298. In Proceedings of the Fourth National Symposium on

- Oceanography and Fisheries, 26-29 April 1993, Rhodes Island. (In Hellenic)
- Crivelli, A.J. and F. Dupont. 1987. Biometrical and biological features of *Alburnus alburnus* x *Rutilus rutilus* natural hybrids from Lake Mikri Prespa, northern Greece. J. Fish Biol. 31:721-733.
- Crivelli, A.J., M. Malakou, G. Catsadorakis and E. Rosecchi. 1996. The Prespa barbel, *Barbus prespensis*, a fish species endemic to the Prespa Lakes (North-Western Greece). Folia. Zool. 45 (Suppl.1): 21-32.
- Daoulas, C. 1981. Contribution to the biology of *Rutilus rutilus* (Bonaparte, 1837), (Pisces, Cyprinidae), in Lake Trichonis (Greece). University of Thessaloniki, Thessaloniki, Hellas, 143 p. Doctorate thesis. (In Hellenic, with English abstract)
- Daoulas, C. and P. Economidis. 1989. Age, growth and feeding of *Barbus albanicus* Steindachner in the Kremasta reservoir, Greece. Arch. Hydrobiol. 114(4):591-601.
- Daoulas, C., T. Koussouris and T. Psarras. 1987. Ecology and possibilities of fisheries management of the artificial lake of Kremasta. Spec. Publ. Natl. Cent. Mar. Res. (Greece), 12:120 p. (In Hellenic)
- Economidis, P.S. 1991. Check list of freshwater fishes of Greece. (Recent status of threats and protection). Hellenic Society for the Protection of Nature, Athens. 48 p.
- Economidis, P.S. and A.I. Sinis. 1982. Les poissons du système des lacs Koronia et Volvi (Macédoine, Grèce). Considérations systématiques et zoogéographiques. Biologia Gallo-Hellenica 9(2):291-317.
- Economou, A.N., C. Daoulas and T. Psarras. 1991a. Growth and morphological development of chub, *Leuciscus cephalus* (L.), during the first year of life. J. Fish Biol. 39:393-408.
- Economou, A.N., C. Daoulas and P. Economidis. 1991b. Observations on the biology of *Leuciscus 'svallize'* in the Kremasta reservoir (Greece). Hydrobiologia 213:99-111.
- Goncalves, J.M.S., L. Bentes, P.G. Lino, J. Ribeiro, A.V.M. Canario and K. Erzini. 1997. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the

- south and south-west coast of Portugal. Fish. Res. 30:253-256.
- Kevrekidis, T., A.K. Kokkinakis and A. Koukouras. 1990. Some aspects of the biology and ecology of *Knipowitschia caucasica* (Teleostei: Gobiidae) in the Evros Delta (North Aegean Sea). Helgol. Meeresunters. 44:173-187.
- Klossa-Kilia, E. 1990. Contribution to the study of the biology of the *Salmo trutta macrostigma* of Acheloos River. University of Patras, Patras, Hellas. Doctorate thesis. 261 p. (In Hellenic)
- Kokkinakis, A. 1992. Comparative study of the biology and dynamics of the fish *Chalcalburnus chalcoides macedonicus* Stephanidis, 1971 (Pisces: Cyprinidae) of the systems Volvi and Vistonis. University of Thessaloniki, Thessaloniki, Hellas. Doctorate thesis. 261 p. (In Hellenic, with English abstract)
- Leonardos, I. 1996. Population dynamics of toothcarp (*Aphanius fasciatus* Nardo, 1827) in the Mesolongi and Etolikon lagoons. University of Thessaloniki, Thessaloniki, Hellas. Doctorate thesis. 198 p. (In Hellenic)
- Neophitou, C. 1987. A study of some autoecological parameters of southern barbel (*Barbus meridionalis* R.) in the Rentina stream, Greece. J. Appl. Ichthyol. 3:24-29.
- Neophitou, C. 1988. Autecology of chub, *Leuciscus cephalus* (L.), in a Greek stream, and the use of the pharyngeal bone in fish predator-prey studies. Aquacult. Fish. Manage. 19:179-190.
- Neophitou, C. 1993a. Ecological study of perch (*Perca fluviatilis* L.) in Lake Doirani. Geot. Scient. Issue. 4(3):38-47.
- Neophitou, C. 1993b. Some biological data on tench (*Tinca tinca* (L.)) in Lake Pamvotida (Greece). Acta Hydrobiol. 35(4):367-379.
- Neophitou, C. and V. Theochari. 1989. Biology of the south European roach (*Rutilus rubilio*) in Lake Pamvotida. Sci. An. Dep. For. Nat. Env. 13:50-110. (In Hellenic, with English abstract)
- Papageorgiou, N. 1977. Age and growth of perch, *Perca fluviatilis* (L.) in the lake of Ag. Vasileios. Thalassographica 1(3):245-265. (In Hellenic)
- Papageorgiou, N. 1979. The length weight relationship, age, growth and reproduction of the roach *Rutilus rutilus* (L.) in Lake Volvi. J. Fish Biol. 14:529-538.
- Papageorgiou, N., C.N. Neophitou and C.G. Vlachos. 1983. The age, growth, and reproduction of brown trout (*Salmo trutta fario*) in the Aspropotamos stream. Acta Hydrobiol. 25/26(3/4):451-467.
- Pauly, D. 1993. Fishbyte section editorial. Naga, ICLARM Q. 16:26.
- Petrakis, G. and K.I. Stergiou. 1995. Weight-length relationships for 33 fish species in Greek waters. Fish. Res. 21:465-469.
- Politou, C-Y. 1993. Biology and dynamics of the fish *Alburnus alburnus* (L., 1758) in lake Koronia. University of Thessaloniki, Thessaloniki, Hellas. Doctorate thesis. 134 p. (In Hellenic, with English abstract)
- Psarras, Th., R. Barbieri-Tseliki and A.N. Economou. 1997. First data on the feeding and biology of reproduction of *Salaria fluviatilis*, p. 261-264. In Proceedings of the Fifth National Symposium on Oceanography and Fisheries, 15-18 April 1997, Kavala, Greece. (In Hellenic, with English abstract)
- Rosecchi, E., A.J. Crivelli and G. Catsadorakis. 1993. The establishment and impact of *Pseudorasbora parva*, an exotic fish species introduced into Lake Mikri Prespa (north-western Greece). Aquat. Cons. Mar. Freshwat. Ecos. 3:223-231.
- Sinis, A. 1981. L'autoécologie de l'espèce endémique *Alosa* (*Caspialosa*) *macedonica* (Vinciguerra) (Pisces: Clupeidae), du lac Volvi. University of Thessaloniki, Thessaloniki, Hellas, 198 p. Doctorate thesis. (In Hellenic, with French abstract)
- Sinis, A. and D. Petridis. 1993. Population structure and reproductive strategy of the whitefish *Coregonus lavaretus* (L.) in two Greek Lakes. Arch. Hydrobiol. 128(4):483-497.
- Sinis, A. and D. Petridis. 1995. Age structure and reproductive pattern of *Chalcalburnus belvica* (Karaman, 1924) in Lake Mikri Prespa (North-western Greece). Isr. J. Zool. 41:569-580.
- Stoumboudi, M.Th., Th. Psarras and R. Barbieri-Tseliki. 1997. Reproductive cycles of *atherina* (*Atherina boyeri* Risso, 1810) from Trichonis Lake (Greece), p. 257-260. In Proceedings of the Fifth National Symposium on Oceanography and Fisheries, 15-18 April 1997, Kavala, Greece. (In Hellenic, with English abstract)
- Tsimenidis, N. 1976. The relationship between fish length and the length of the operculum for the carp in Lake Vistonis. Thalassographica 1(1):53-63.
- Valoukas, V.A. and P.S. Economidis. 1996. Growth, population composition and reproduction of Bream *Abramis brama* (L.) in Lake Volvi, Macedonia, Greece. Ecol. Freshwat. Fish. 5:108-115.

P.K. KLEANTHIDIS, A.I. SINIS and K.I. STERGIOU are from the Laboratory of Ichthyology, Department of Zoology, P.O. Box 134, Faculty of Sciences, Aristotle University of Thessaloniki, 54006, Thessaloniki, Hellas.